arXiv:0805.0052v1 [math.NT] 1 May 2008

On a variance for twins of $k-$free numbers in arithmetic progressions [*]

Zaizhao MENG

In this paper, we give a new upper bound of Barban-Davenport-Halberstam type for twins of $k-$free numbers in arithmetic progressions.

1 Introduction

In recent years, Brüdern and others [1,2] have made a breakthrough in the circle method. They give successful treatment, through the circle method, of binary additive problems involving $k-$free numbers. Their results depend upon the variance for $k-$free numbers (or twins of $k-$free numbers) in arithmetic progressions.

In this paper, we give new results on the variance for twins of $k-$free numbers in arithmetic progressions. Such result is analogous to the Barban-Davenport-Halberstam theorem for the primes in arithmetic progressions.

Let $\mu_k(n)$ be the characteristic function of the $k-$free numbers,

$$\mu_k(n)=\sum_{d^k|n}\mu(d), \tag{1.1}$$

where $\mu(n)$ is the Möbius function.

Let

$$A_k(x;q,a)=\sum_{\substack{n\le x\\ n\equiv a(mod\ q)}}\mu_k(n)\mu_k(n+1) \tag{1.2}$$

---

[*] 2000 Mathematics Subject Classification: 11N25.

$$g(q,a)=\sum_{\substack{u,v=1\\(u^k,q)|a,(v^k,q)|a+1}}^{\infty}\frac{\mu(uv)}{u^kv^k}(q,u^kv^k) \tag{1.3}$$

By Lemma 3.1 in [2], we have

$$A_k(x;q,a)=q^{-1}g(q,a)x+O(x^{\frac{2}{k+1}+\varepsilon}).$$

We consider the variance

$$Y_k(x,Q)=\sum_{q\le Q}\sum_{a=1}^{q}\mid A_k(x;q,a)-q^{-1}g(q,a)x\mid^2 . \tag{1.4}$$

In [2], Brüdern, Perelli and Wooley obtained

$$Y_k(x,Q)\ll x^{\frac{2}{k}+\varepsilon}Q^{2-\frac{2}{k}}+x^{\frac{4}{k+1}+\varepsilon},\quad 1<Q\le x. \tag{1.5}$$

In this paper, we obtain the following

THEOREM. Suppose that $1<Q\le x$ and that k is an integer with $k\ge 2$. Then

$$Y_k(x,Q)\ll x^{\frac{1}{k}+\varepsilon}Q^{2-\frac{1}{k}-\varepsilon}+x^{1+\frac{2}{k}}\log Q+x^{\frac{3}{2}+\frac{1}{2k}+\varepsilon}.$$

Actually, the first term is $Q^2(\frac{x}{Q})^{\frac{1}{k}+\varepsilon}$. For $k>2$, this result is superior to (1.5) when $Q\gg x^{\frac{3}{4}}$.

We have

$$Y_k(x,Q)=S_1(x,Q)-2xS_2(x,Q)+x^2S_3(x,Q) \tag{1.6}$$

where

$$S_1(x,Q)=\sum_{q\le Q}\sum_{a=1}^{q}A_k^2(x;q,a), \tag{1.7}$$

$$S_2(x,Q)=\sum_{q\le Q}\sum_{a=1}^{q}A_k(x;q,a)q^{-1}g(q,a), \tag{1.8}$$

$$S_3(x,Q)=\sum_{q\le Q}\sum_{a=1}^{q}q^{-2}g(q,a)^2. \tag{1.9}$$

We notice that the function $g(q,a)$ does not depend only on q and (a,q), unlike most sequences investigated before.

In section 2-3, we discuss $S_2(x,Q)$ and $S_3(x,Q)$, these depend on the solutions of the corresponding congruences. In section 4-7, we discuss $S_1(x,Q)$ by the Hardy-Littlewood method on the line of Vaughan[9]. In section 8, we discuss the singular series $\mathfrak{S}(n)$. In section 9-10, we use the Hurwitz Zeta function to discuss the generating function of $\mathfrak{S}(uv)$. In section 11, we use a trick to avoid the difficulty of calculating some constants and then we obtain the Theorem.

Notation: Let $k > 1$ denote a positive integer. Throughout, ε is a sufficiently small positive number, the implicit constants in Vinogradov's notation $\ll$, and in Landau's O-notation, will depend at most on k, ε unless it is pointed out depend upon the corresponding parameters. $n \equiv a(mod\ q)$ may be written as $n \equiv a(q)$. The greatest common divisor and the least common multiple of integers a, b are denoted by (a,b) and $[a,b]$ respectively; $\mu(n)$ denotes the Möbius function and $\tau(n)$ denotes the divisor function; $[w]$ denotes the integer part of w and $e(\alpha) = \exp(2\pi i\alpha)$; $\sum_{a=1}^{q}{}'$ means $(a,q) = 1$. The letter p denotes a prime number, and write $p^t \parallel n$ when $p^t \mid n$ but $p^{t+1} \dagger n$. Let x denote a sufficiently large real number and Q be a positive real number with $1 < Q \le x$.

2 The formula for $S_2(x,Q)$

For fixed positive integers u, v, q, r, s, let

$$J_1 = \sum_{\substack{a=1 \\ (u^k,q)|a,(v^k,q)|a+1}}^{q} \sum_{\substack{n\le x, n\equiv a(mod\ q) \\ r^k|n, s^k|n+1}} 1, \quad s_0 = (\frac{(u^k,q)}{(r^k,u^k,q)}, \frac{r^k}{(r^k,q)}). \tag{2.1}$$

LEMMA 2.1. We have

$$J_1 = \frac{(r^k,u^k,q)s_0}{r^k s^k (u^k,q)(v^k,q)}(s^k, \frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)})x + O(1),$$

provided that

$$(\frac{(u^k,q)(r^k,q)}{(r^k,u^k,q)}, (v^k,q)) = (\frac{(u^k,q)}{(r^k,u^k,q)s_0}, s^k) = (r,s) = (q,u,v) = 1. \tag{2.2}$$

Proof. We use an idea of [4]. Let $n = r^k b,\ a = (u^k, q)c$, then

$$J_1 = \sum_{\substack{c=1 \\ (v^k,q)|(u^k,q)c+1}}^{q(u^k,q)^{-1}} \sum_{\substack{r^k b \le x \\ r^k b \equiv (u^k,q)c(mod\ q), s^k | r^k b+1}} 1, \quad (r, s) = (q, u, v) = 1. \tag{2.3}$$

Now,

$\frac{r^k}{(r^k,q)} b \equiv \frac{(u^k,q)c}{(r^k,q)} (mod\ \frac{q}{(r^k,q)}),\quad (r^k, q) \mid (u^k, q)c.$

Since $\frac{(u^k,q)(r^k,q)}{(r^k,u^k,q)} = [(u^k, q), (r^k, q)] \mid q$, we write $c = \frac{(r^k,q)}{(r^k,u^k,q)} d$, then

$J_1 = \#\{1 \le b \le xr^{-k},\ 1 \le d \le \frac{q(r^k,u^k,q)}{(u^k,q)(r^k,q)} :\ (v^k, q) \mid (u^k, q)\frac{(r^k,q)}{(r^k,u^k,q)} d + 1,$

$\frac{r^k}{(r^k,q)} b \equiv \frac{(u^k,q)}{(r^k,q)} \frac{(r^k,q)}{(r^k,u^k,q)} d(mod\ \frac{q}{(r^k,q)}),\ s^k \mid r^k b + 1.\}$

We have

$$\frac{r^k}{(r^k, q)} b \equiv \frac{(u^k, q)d}{(r^k, u^k, q)} (mod\ \frac{q}{(r^k, q)}),$$

hence

$\frac{r^k}{(r^k,q)} \frac{(r^k,u^k,q)}{(u^k,q)} b \equiv d(mod\ \frac{q}{(r^k,q)} \frac{(r^k,u^k,q)}{(u^k,q)}),$

and

$$\frac{(u^k, q)}{(r^k, u^k, q)} \mid \frac{r^k}{(r^k, q)} b. \tag{2.4}$$

Write $b = \frac{(u^k,q)}{(r^k,u^k,q)s_0} b_1$, then

$J_1 = \#\{1 \le \frac{(u^k,q)}{(r^k,u^k,q)s_0} b_1 \le xr^{-k},\ 1 \le d \le \frac{q(r^k,u^k,q)}{(u^k,q)(r^k,q)} :\ (v^k, q) \mid (u^k, q)\frac{(r^k,q)}{(r^k,u^k,q)} d + 1,$

$\frac{r^k}{(r^k,q)s_0} b_1 \equiv d(mod\ \frac{q(r^k,u^k,q)}{(r^k,q)(u^k,q)}),\ s^k \mid r^k \frac{(u^k,q)}{(r^k,u^k,q)s_0} b_1 + 1.\}$

We have

$$\frac{(u^k, q)(r^k, q)}{(r^k, u^k, q)} d \equiv -1(mod\ (v^k, q)), \tag{2.5}$$

and

$$(\frac{(u^k, q)(r^k, q)}{(r^k, u^k, q)}, (v^k, q)) = 1, \tag{2.6}$$

$$(v^k, q) \mid \frac{q(r^k, u^k, q)}{(u^k, q)(r^k, q)}, \tag{2.7}$$

$$(\frac{(u^k,q)}{(r^k,u^k,q)s_0},s^k)=1. \tag{2.8}$$

We write

$d=d_1+m(v^k,q),\ 1\le d_1\le (v^k,q),\ 0\le m<\frac{q(r^k,u^k,q)}{(v^k,q)(u^k,q)(r^k,q)},$

then

$$\frac{r^k}{(r^k,q)s_0}b_1\equiv d_1+m(v^k,q)(mod\ \frac{q(r^k,u^k,q)}{(r^k,q)(u^k,q)}) \tag{2.9}$$

$$\frac{r^k}{(r^k,q)s_0}b_1\equiv d_1(mod\ (v^k,q)). \tag{2.10}$$

If a prime $p\mid (v,q)$, then $p\mid v,\ p\mid q, p\dagger u$, again if $p\mid \frac{r^k}{(r^k,q)s_0}$, then $p\mid r$, hence,by (2.6) $p\mid (\frac{(u^k,q)(r^k,q)}{(r^k,u^k,q)},(v^k,q))=1$, this is a contradiction.

We deduce that

$$((v^k,q),\frac{r^k}{(r^k,q)s_0})=1. \tag{2.11}$$

Now the congruence (2.10) has a unique solution modulus (v^k,q).

Let $b_1=b_0+t(v^k,q),\ 1\le b_0\le (v^k,q)$, by (2.7) and (2.9)

$$\frac{r^k(v^k,q)}{(r^k,q)s_0}t\equiv d_1-\frac{r^k}{(r^k,q)s_0}b_0+m(v^k,q)(mod\ \frac{q(r^k,u^k,q)}{(r^k,q)(u^k,q)}). \tag{2.12}$$

Also, we have

$$\frac{r^k(u^k,q)}{(r^k,u^k,q)s_0}(v^k,q)t\equiv -1-\frac{r^k(u^k,q)}{(r^k,u^k,q)s_0}b_0\ (mod\ s^k). \tag{2.13}$$

By (2.7), (2.10) and (2.12),

$$\frac{r^k}{(r^k,q)s_0}t\equiv (d_1-\frac{r^k}{(r^k,q)s_0}b_0)(v^k,q)^{-1}+m(mod\ \frac{q(r^k,u^k,q)}{(r^k,q)(u^k,q)(v^k,q)}), \tag{2.14}$$

and

$$d_1\equiv \frac{r^k}{(r^k,q)s_0}b_0\ (mod\ (v^k,q)). \tag{2.15}$$

By (2.5),

$$\frac{(u^k,q)(r^k,q)}{(r^k,u^k,q)}d_1 \equiv -1(mod\ (v^k,q)). \tag{2.16}$$

By (2.14), we only want to count the number of t satisfying (2.13) and

$$\frac{(u^k,q)}{(r^k,u^k,q)s_0}(b_0+t(v^k,q)) \leq xr^{-k},$$

that is

$$t \leq \frac{x(r^k,u^k,q)s_0}{r^k(u^k,q)(v^k,q)} - \frac{b_0}{(v^k,q)}. \tag{2.17}$$

By (2.5), (2.15) and (2.16),

$$\frac{(u^k,q)r^k}{(r^k,u^k,q)s_0}b_0 \equiv -1(mod\ (v^k,q)). \tag{2.18}$$

Consequently, by(2.13) and (2.18),

$$((v^k,q),s^k) \mid 1+\frac{(u^k,q)r^k}{(r^k,u^k,q)s_0}b_0. \tag{2.19}$$

By (2.3), (2.8) and (2.19), we have

$$(s^k,\frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)s_0}) \mid 1+\frac{(u^k,q)r^k}{(r^k,u^k,q)s_0}b_0.$$

By (2.13) and $(s,s_0)=1$, we obtain

$\frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)}(s^k,\frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)})^{-1}t$
$\equiv (-s_0-\frac{r^k(u^k,q)}{(r^k,u^k,q)}b_0)(s^k,\frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)})^{-1}\ (mod\ s^k(s^k,\frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)})^{-1}).$

Therefore, the number of t is

$$\frac{x(r^k,u^k,q)s_0}{r^k(u^k,q)(v^k,q)}(s^k(s^k,\frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)})^{-1})^{-1}+O(1),$$

that is

$$J_1=\frac{(r^k,u^k,q)s_0}{r^ks^k(u^k,q)(v^k,q)}(s^k,\frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)})x+O(1).$$

The lemma follows.

Now,we give the formula for $S_2(x,Q)$.

$$S_2(x,Q) = \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \frac{\mu(uv)}{u^k v^k}(q,u^k v^k) \sum_{\substack{a=1 \\ (u^k,q)|a,(v^k,q)|a+1}}^{q} \sum_{\substack{n\le x \\ n\equiv a(mod\ q)}} \mu_k(n)\mu_k(n+1)$$

$$= \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \frac{\mu(uv)}{u^k v^k}(q,u^k v^k) \sum_{\substack{r^k\le x \\ s^k\le x+1}} \mu(r)\mu(s)J_1.$$

By Lemma 2.1,

$$S_2(x,Q) = S_{2A}(x,Q) + S_{2B}(x,Q), \tag{2.20}$$

where

$$S_{2A}(x,Q) = \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \frac{\mu(uv)}{u^k v^k}(q,u^k v^k) \sum_{\substack{r^k\le x, s^k\le x+1 \\ (2.1),(2.2)}} \mu(r)\mu(s)\frac{(r^k,u^k,q)s_0}{r^k s^k (u^k,q)(v^k,q)}(s^k, \frac{r^k(u^k,q)(v^k,q)}{(r^k,u^k,q)})x,$$

$$S_{2B}(x,Q) \ll \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \frac{\mu(uv)}{u^k v^k}(q,u^k v^k) \sum_{\substack{r^k\le x \\ s^k\le x+1}} 1,$$

and $\sum\limits_{\substack{r^k\le x, s^k\le x+1 \\ (2.1),(2.2)}}$ means u,v,r,s,q satisfy (2.1) and (2.2).

We give upper bound for $S_{2B}(x,Q)$ first.

$$S_{2B}(x,Q) \ll x^{\frac{2}{k}} \sum_{m=1}^{\infty} \frac{\mu^2(m)\tau(m)}{m^k} \sum_{q\le Q} q^{-1}(q,m^k)$$

$$\ll x^{\frac{2}{k}} \sum_{m=1}^{\infty} \frac{\mu^2(m)\tau(m)}{m^k} \sum_{t|m^k} \sum_{\substack{q\le Q \\ (q,m^k)=t}} q^{-1}(q,m^k) \ll x^{\frac{2}{k}} \sum_{m=1}^{\infty} \frac{\mu^2(m)\tau(m)}{m^k} \sum_{t|m^k} \sum_{qt\le Q} q^{-1},$$

hence

$$S_{2B}(x,Q) \ll x^{\frac{2}{k}} \log Q. \tag{2.21}$$

By $(\frac{(u^k,q)}{(r^k,u^k,q)}, \frac{r^k}{(r^k,q)}\frac{(r^k,q)}{(r^k,u^k,q)}) = (\frac{(u^k,q)}{(r^k,u^k,q)}, \frac{r^k}{(r^k,u^k,q)}) = 1$, we have

$$s_0 = (\frac{(u^k,q)}{(r^k,u^k,q)}, \frac{r^k}{(r^k,q)}) = 1. \tag{2.22}$$

Since $(u,v)=1$, the conditions (2.2) and (2.22) are equivalent to

$$(r,v,q) = (s,u,q) = (s,r) = 1. \tag{2.23}$$

By (2.2),

$$S_{2A}(x,Q) = x\sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \frac{\mu(uv)}{u^k v^k} \sum_{\substack{r^k\le x, s^k\le x+1 \\ (2.23)}} \mu(r)\mu(s)r^{-k}s^{-k}(s^k,v^k,q)(r^k,u^k,q). \tag{2.24}$$

The summations of r, s can be completed to ∞.

$$S_{2A}(x,Q) = c_1(Q)x + S_{2A_0}, \tag{2.25}$$

where

$$c_1(Q) = \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \frac{\mu(uv)}{u^k v^k} \sum_{\substack{r,s=1\\(2.23)}}^{\infty} \mu(r)\mu(s) r^{-k} s^{-k} (s^k, v^k, q)(r^k, u^k, q), \tag{2.26}$$

and

$$S_{2A_0} \ll x \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \Big(\sum_{\substack{r^k > x\\ s^k \le x+1, (r,s)=1}} r^{-k} s^{-k} (s^k, v^k, q)(r^k, u^k, q) +$$

$$\sum_{\substack{r\ge 1\\ s^k > x+1, (r,s)=1}} r^{-k} s^{-k} (s^k, v^k, q)(r^k, u^k, q)\Big) = S_{2Aa} + S_{2Ab}, \text{ say.}$$

$$S_{2Aa} \ll x \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \sum_{s^k \le x+1} s^{-k} (s^k, v^k, q) \sum_{t|u} \sum_{\substack{r^k > x\\ (r,u)=t, (r,s)=1}} r^{-k} (t^k, q)$$

$$\ll x \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \sum_{s^k \le x+1} s^{-k} (s^k, v^k, q) \sum_{t|u, (t,s)=1} (t^k, q) t^{-k} \sum_{t^k r^k > x} r^{-k}.$$

Now

$$\sum_{t|u,(t,s)=1} (t^k, q) t^{-k} \sum_{t^k r^k > x} r^{-k}$$

$$\ll \sum_{\substack{t|u,(t,s)=1\\ t^k \le x}} (t^k, q) t^{-k} (x^{\frac{1}{k}} t^{-1})^{1-k} + \sum_{\substack{t|u\\ t^k > x}} (t^k, q) t^{-k} \ll x^{\frac{1}{k}-1} \sum_{\substack{t|u,(t,s)=1\\ t^k \le x}} (t^k, q) t^{-1} + \sum_{\substack{t|u,(t,s)=1\\ t^k > x}} (t^k, q) t^{-k}.$$

Hence

$$S_{2Aa} \ll x \sum_{q\le Q} q^{-1} \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \sum_{s^k \le x+1} s^{-k} (s^k, v^k, q) \Big(x^{\frac{1}{k}-1} \sum_{\substack{t|u,(t,s)=1\\ t^k \le x}} (t^k, q) t^{-1} + \sum_{\substack{t|u,(t,s)=1\\ t^k > x}} (t^k, q) t^{-k}\Big)$$

$$\ll S_{2Aa1} + S_{2Aa2}, \text{ say.}$$

Firstly, we estimate S_{2Aa1},

$$S_{2Aa1} \ll x^{\frac{1}{k}} \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \sum_{s^k \le x+1} s^{-k} \sum_{\substack{t|u,(t,s)=1\\ t^k \le x}} t^{-1} \sum_{q\le Q} q^{-1} (s^k, v^k, q)(t^k, q)$$

$$\ll x^{\frac{1}{k}} \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \sum_{s^k \le x+1} s^{-k} \sum_{\substack{t|u\\ t^k \le x}} t^{-1} \sum_{q\le Q} q^{-1} (s^k t^k, q)$$

$$\ll x^{\frac{1}{k}} \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \sum_{s^k \le x+1} s^{-k} \sum_{\substack{t|u\\ t^k \le x}} t^{-1} \sum_{j|s^k t^k} \sum_{qj\le Q} q^{-1} \ll x^{\frac{1}{k}} \log Q,$$

hence

$$S_{2Aa1} \ll x^{\frac{1}{k}} \log Q. \tag{2.27}$$

Secondly, as the estimate of S_{2Aa1}, we have

$$S_{2Aa2} \ll x \sum_{q \le Q} q^{-1} \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \sum_{s^k \le x+1} s^{-k} (s^k, v^k, q) \sum_{\substack{t|u,(t,s)=1 \\ t^k > x}} (t^k, q) t^{-k}$$

$$\ll x \sum_{u,v=1}^{\infty} \tfrac{1}{u^k v^k} \sum_{s^k \le x+1} s^{-k} \sum_{\substack{t|u,(t,s)=1 \\ t^k > x}} \sum_{q \le Q} q^{-1} (s^k, v^k, q)(t^k, q) t^{-k}$$

$$\ll x \log Q \sum_{v=1}^{\infty} \tfrac{1}{v^k} \sum_{s^k \le x+1} s^{-k} \tau(s^k) (x^{\frac{1}{k}})^{1-2k+\varepsilon},$$

hence

$$S_{2Aa2} \ll x^{\frac{1}{k}-1+\varepsilon} \log Q. \tag{2.28}$$

By (2.27) and (2.28),

$$S_{2Aa} \ll x^{\frac{1}{k}} \log Q + x^{\frac{1}{k}-1+\varepsilon} \log Q. \tag{2.29}$$

In the same manner we have

$$S_{2Ab} \ll x^{\frac{1}{k}} \log Q + x^{\frac{1}{k}-1+\varepsilon} \log Q. \tag{2.30}$$

By (2.29) and (2.30),

$$S_{2A_0} \ll x^{\frac{1}{k}} \log Q + x^{\frac{1}{k}-1+\varepsilon} \log Q. \tag{2.31}$$

By (2.20), (2.21), (2.25) and (2.31), we obtain

$$S_2(x, Q) = c_1(Q) x + O(x^{\frac{2}{k}} \log Q), \tag{2.32}$$

where $c_1(Q)$ is given by (2.26).

3 The formula for $S_3(x, Q)$

By (1.3) and (1.9),

$$S_3(x, Q) = \sum_{q \le Q} \sum_{a=1}^{q} q^{-2} \sum_{\substack{u,v=1 \\ (u^k,q)|a,(v^k,q)|a+1}}^{\infty} \frac{\mu(uv)}{u^k v^k} (q, u^k v^k) \sum_{\substack{r,s=1 \\ (r^k,q)|a,(s^k,q)|a+1}}^{\infty} \frac{\mu(rs)}{r^k s^k} (q, r^k s^k). \tag{3.1}$$

Let

$$\Delta = \#\{1 \le a \le q : \ (u^k, q) \mid a, \ (r^k, q) \mid a, \ (v^k, q) \mid a+1, \ (s^k, q) \mid a+1\}. \tag{3.2}$$

Then

$$(u, s, q) = (v, r, q) = 1,$$

and

$$\Delta = \#\{1 \le m \le \frac{q}{[(u^k, q), (r^k, q)]} : \ m[(u^k, q), (r^k, q)] + 1 \equiv 0 (mod\ [(v^k, q), (s^k, q)])\},$$

therefore,

$$([(u^k, q), (r^k, q)], [(v^k, q), (s^k, q)]) = 1,$$

and

$$\Delta = \frac{q}{[(u^k, q), (r^k, q)][(v^k, q), (s^k, q)]} = \frac{q(r^k, u^k, q)(s^k, v^k, q)}{(u^k, q)(r^k, q)(v^k, q)(s^k, q)}. \tag{3.3}$$

Hence, by (3.1), (3.2) and (3.3),

$$S_3(x, Q) = c_1(Q), \tag{3.4}$$

where $c_1(Q)$ is given by (2.26).

4 Lemmas for $S_1(x, Q)$

As Vaughan had done in [9], our proof of the theorem uses the Hardy-Littlewood method(see [8]) and depends heavily on [2] where the bounds for

$$S(\alpha) = S_k(\alpha) = \sum_{n \le x} \mu_k(n)\mu_k(n+1)e(n\alpha), \tag{4.1}$$

and related expressions, are obtained.

Let $R = x^{1/2+\tau}$, where $\tau = \tau_k$ is a sufficiently small positive number($2\varepsilon < \tau < \frac{1}{10k}$), and let $\mathfrak{M}$ denote the union of the intervals

$$\mathfrak{M}(q, a) = \{\alpha :\mid q\alpha - a \mid \le R^{-1}\}, \tag{4.2}$$

with $1 \leq a \leq q \leq x/R$ and $(q,a)=1$, and $\mathscr{M} = (R^{-1}, 1+R^{-1}]\backslash\mathfrak{M}$.

LEMMA 4.1. We have

$$\int_{\mathscr{M}} \mid S(\alpha) \mid^2 d\alpha \ll x^{\frac{1}{k}+\varepsilon} R^{1-\frac{1}{k}} + x^{2+\varepsilon} R^{\frac{2}{k}-3} + x^{\frac{4}{k+1}+1+\varepsilon} R^{-2}.$$

Proof. This is Theorem 2 of [2] by taking $Q = x/R$, we note that $\mathfrak{M} \subseteq (R^{-1}, 1+R^{-1}]$, the interval $(Q^{-1}, 1+Q^{-1}]$ of [2] may be replaced by $(R^{-1}, 1+R^{-1}]$.

Let

$$K(\alpha) = \sum_{u \leq Q} \sum_{v \leq x/u} e(uv\alpha). \tag{4.3}$$

When $Q \leq \sqrt{x}$ define

$$K_q(\alpha) = \sum_{u \leq Q/q} \sum_{v \leq x/qu} e(quv\alpha), \tag{4.4}$$

$$H_q(\alpha) = \begin{cases} \sum\limits_{\substack{u \leq Q \\ q \dagger u}} \sum\limits_{v \leq x/u} e(uv\alpha), & (q > 1), \\ 0, & (q = 1). \end{cases} \tag{4.5}$$

When $Q > \sqrt{x}$ define

$$K_q(\alpha) = \sum_{u \leq \sqrt{x}/q} \Big(\sum_{v \leq x/qu} e(quv\alpha) + \sum_{\sqrt{x} < v \leq \min(Q, x/qu)} e(quv\alpha) \Big), \tag{4.6}$$

$$H_q(\alpha) = \begin{cases} \sum\limits_{\substack{u \leq \sqrt{x} \\ q \dagger u}} \Big(\sum\limits_{v \leq x/u} e(uv\alpha) + \sum\limits_{\sqrt{x} < v \leq \min(Q, x/u)} e(uv\alpha) \Big), & (q > 1), \\ 0, & (q = 1). \end{cases} \tag{4.7}$$

We have the following results on these functions(see the section 2 of Vaughan [9]).

LEMMA 4.2. Suppose that $Q \leq x$ and $q \in N$. Then

$$K(\alpha) = K_q(\alpha) + H_q(\alpha).$$

Proof. This is Lemma 2.9 of [9].

LEMMA 4.3. Suppose that $(a,q) = 1$ and $\mid q\alpha - a \mid \leq q^{-1}$. Then

$$K(\alpha) \ll (xq^{-1} + q) \log x.$$

Proof. This is Lemma 2.10 of [9].

LEMMA 4.4. Suppose that $(a,q)=1, q\in N$ and $\alpha = a/q+\beta$. Then

$$K_q(\alpha)=K_q(\beta)\ll \sum_{u\le\sqrt{x}/q}\frac{x}{qu+x\|qu\beta\|}.$$

Moreover, if $|\beta|\le 1/(2\sqrt{x})$, then

$$K_q(\alpha)\ll\frac{x\log x}{q+qx\mid\beta\mid}.$$

Proof. This is Lemma 2.11 of [9].

LEMMA 4.5. Suppose that $(a,q)=1, q\in N$ and $|q\alpha-a|\le 1/(2\sqrt{x})$. Then

$$H_q(\alpha)\ll(\min(Q,\sqrt{x})+q)\log x.$$

Proof. This is Lemma 2.12 of [9].

By Dirichlet's theorem on diophantine approximation, Lemma 4.1 and Lemma 4.3,

$$\sup_{\mathscr{M}}\mid K(\alpha)\mid\ll R\log x \tag{4.8}$$

and

$\int_{\mathscr{M}}\mid S(\alpha)\mid^2 K(\alpha)d\alpha\ll x^{\frac{1}{k}+\varepsilon}R^{2-\frac{1}{k}}+x^{2+\varepsilon}R^{\frac{2}{k}-2}+x^{\frac{4}{k+1}+1+\varepsilon}R^{-1}$

$$\ll x^{1+\frac{1}{k}-\frac{\tau}{2}}+x^{\frac{1}{2}+\frac{4}{k+1}-\frac{\tau}{2}}. \tag{4.9}$$

By (1.2) and (1.7),

$S_1(x,Q)=\sum\limits_{q\le Q}\sum\limits_{\substack{m,n\le x\\ m\equiv n(mod\ q)}}\mu_k(n)\mu_k(n+1)\mu_k(m)\mu_k(m+1)$
$=2\sum\limits_{q\le Q}\sum\limits_{\substack{m<n\le x\\ m\equiv n(mod\ q)}}\mu_k(n)\mu_k(n+1)\mu_k(m)\mu_k(m+1)+[Q]\sum\limits_{n\le x}\mu_k(n)\mu_k(n+1).$

Hence

$$S_1(x,Q)=2S_A(x,Q)+[Q]\sum_{n\le x}\mu_k(n)\mu_k(n+1) \tag{4.10}$$

where

$$S_A(x,Q)=\sum_{q\le Q}\sum_{\substack{m<n\le x\\ m\equiv n(mod\ q)}}\mu_k(n)\mu_k(n+1)\mu_k(m)\mu_k(m+1). \tag{4.11}$$

By (4.1), (4.3) and (4.11),

$$S_A(x,Q)=\int_{R^{-1}}^{1+R^{-1}} \mid S(\alpha)\mid^2 K(\alpha)d\alpha=\int_{\mathfrak{M}} \mid S(\alpha)\mid^2 K(\alpha)d\alpha+\int_{\mathscr{M}} \mid S(\alpha)\mid^2 K(\alpha)d\alpha. \tag{4.12}$$

By (4.9) and (4.12),

$$S_A(x,Q)=\int_{\mathfrak{M}} \mid S(\alpha)\mid^2 K(\alpha)d\alpha+O(x^{1+\frac{1}{k}-\frac{\tau}{2}}+x^{\frac{1}{2}+\frac{4}{k+1}-\frac{\tau}{2}}). \tag{4.13}$$

By (1.3)

$$g(q,a)=\sum_{n=1}^{\infty}\mu(n)\frac{(n^k,q)}{n^k}\psi_k(n;q,a), \tag{4.14}$$

where $\psi_k(n;q,a)$ denotes the number of pairs u,v of natural numbers with $uv=n,\ (u,v)=1,\ (u^k,q)\mid a,\ (v^k,q)\mid a+1$. For fixed $a,\ q$, the function $\psi_k(n;q,a)$ is multiplicative in n, hence

$$g(q,a)=\prod_p(1-\frac{(p^k,q)}{p^k}\psi_k(p;q,a)).$$

Let

$$f(q)=\prod_{p\mid q}(1-\frac{2}{p^k})^{-1},\quad h(q,a)=\prod_{p\mid q}(1-\frac{(p^k,q)}{p^k}\psi_k(p;q,a)), \tag{4.15}$$

then, by (3.8) of [2],

$$g(q,a)=\varrho f(q)h(q,a), \tag{4.16}$$

where

$$\varrho=\prod_p(1-\frac{2}{p^k}). \tag{4.17}$$

By (3.9) and (3.10) of [2], we have

$$h(q,a)=\prod_{\substack{p\mid q\\ (p^k,q)\mid a(a+1)}}(1-\frac{(p^k,q)}{p^k}), \tag{4.18}$$

and for co-prime natural numbers q_1,q_2

$$h(q_1q_2,a)=h(q_1,a)h(q_2,a), \tag{4.19}$$

$h(q,a)$ is a multiplicative function of q.

We need the following sums of Gaussian type

$$G(q,a)=\sum_{b=1}^{q} g(q,b)e(\frac{ab}{q}),\ \ H(q,a)=\sum_{b=1}^{q} h(q,b)e(\frac{ab}{q}), \tag{4.20}$$

which by (4.16) are related by

$$G(q,a)=\varrho f(q)H(q,a). \tag{4.21}$$

As in [2], we introduce the following function

$$H(q)=\sum_{\substack{a=1\\(a,q)=1}}^{q} \mid H(q,a)\mid^2 . \tag{4.22}$$

We define

$$I(\beta)=\sum_{n\leq x} e(\beta n), \quad S^*(\alpha)=S^*(\alpha;q,a)=q^{-1}G(q,a)I(\alpha-\frac{a}{q}), \tag{4.23}$$

$$\Delta(\alpha)=\Delta(\alpha;q,a)=S(\alpha)-S^*(\alpha). \tag{4.24}$$

Let

$$\mathfrak{M}(q,a;T)=\{\alpha:\mid q\alpha-a\mid\leq T/x\}, \tag{4.25}$$

and $\mathfrak{M}(T)$ denote the union of the intervals $\mathfrak{M}(q,a;T)$ with $1\leq a\leq q\leq T$ and $(q,a)=1$, then

$$\mathfrak{M}(2T)\backslash\mathfrak{M}(T)=\bigcup_{\substack{T<q\leq 2T\\ a\leq q,(a,q)=1}}\{\alpha:\mid q\alpha-a\mid\leq T/x\}\bigcup_{\substack{q\leq T\\ a\leq q,(a,q)=1}}\{\alpha:T/x<\mid q\alpha-a\mid\leq 2T/x\}.$$

and

$$\mathfrak{M}=\mathfrak{M}(x/R)\subseteqq\mathfrak{M}(1)\bigcup_{1\leq 2^j\leq x/R}\mathfrak{M}(2^{j+1})\backslash\mathfrak{M}(2^j). \tag{4.26}$$

Hence, as the proof of Lemma 4.2 in [1],

$$\int_{\mathfrak{M}}\mid S^*(\alpha)\Delta(\alpha)\mid d\alpha\ll\log x\max_{1\leq T\leq x/R}(U_1^*(T))^{1/2}(U_2^*(T))^{1/2}+\int_{\mathfrak{M}(1)}\mid S^*(\alpha)\Delta(\alpha)\mid d\alpha,$$

(4.27)

where

$$U_1^*(T) = \int_{\mathfrak{M}(2T)\backslash\mathfrak{M}(T)} \mid S^*(\alpha) \mid^2 d\alpha,\ U_2^*(T) = \int_{\mathfrak{M}(2T)} \mid \Delta(\alpha) \mid^2 d\alpha. \tag{4.28}$$

LEMMA 4.6. Suppose that $1 \leq T \leq \frac{1}{2}\sqrt{x}$. Then

$$\int_{\mathfrak{M}(T)} \mid S(\alpha) - S^*(\alpha) \mid^2 d\alpha \ll T^{3-\frac{2}{k}} x^{\frac{2}{k}-1+\varepsilon} + x^{\frac{4}{k+1}-1+\varepsilon} T^2.$$

Proof. This is Lemma 5.1 of [2].

LEMMA 4.7. Suppose that $1 \leq 2T \leq \frac{1}{2}\sqrt{x}$. Then

$$\int_{\mathfrak{M}(2T)\backslash\mathfrak{M}(T)} \mid S^*(\alpha) \mid^2 d\alpha \ll x T^{\frac{1}{k}-1+\varepsilon}.$$

Proof. This is Lemma 5.2 of [2].

LEMMA 4.8. We have

$$\int_{\mathfrak{M}} \mid S(\alpha) - S^*(\alpha) \mid^2 d\alpha \ll x^{2+\varepsilon} R^{\frac{2}{k}-3} + x^{\frac{4}{k+1}+1+\varepsilon} R^{-2}.$$

Proof. By taking $T = x/R$ in Lemma 4.6, the lemma follows.

LEMMA 4.9. We have

$$\sum_{q \leq x/R} \sideset{}{'}\sum_{a=1}^{q} \int_{\mathfrak{M}(q,a)} \mid \Delta(\alpha; q, a) \mid^2 H_q(\alpha) d\alpha \ll x^{\frac{5}{2}+\varepsilon} R^{\frac{2}{k}-3} + x^{\frac{3}{2}+\frac{4}{k+1}+\varepsilon} R^{-2}.$$

Proof. By Lemma 4.5 and Lemma 4.6, the lemma follows.

LEMMA 4.10. We have

$$\int_{\mathfrak{M}(1)} \mid S^*(\alpha)\Delta(\alpha) \mid d\alpha \ll x^{\frac{2}{k+1}+\varepsilon}.$$

Proof. By Cauchy-Schwarz inequality and Lemma 4.6

$\int_{\mathfrak{M}(1)} \mid S^*(\alpha)\Delta(\alpha) \mid d\alpha \ll \int_{1-1/x}^{1+1/x} \mid S^*(\alpha)\Delta(\alpha) \mid d\alpha$

$\ll \int_0^{x^{-1}} \mid S^*(\alpha)\Delta(\alpha) \mid d\alpha \ll (\int_0^{x^{-1}} \mid S^*(\alpha) \mid^2 d\alpha)^{1/2} (\int_0^{x^{-1}} \mid \Delta(\alpha) \mid^2 d\alpha)^{1/2}$

$\ll x^{\frac{1}{2}} (x^{\frac{2}{k}-1+\varepsilon} + x^{\frac{4}{k+1}-1+\varepsilon})^{1/2} \ll x^{\frac{2}{k+1}+\varepsilon}$, the lemma follows.

By (4.28), Lemma 4.6 and Lemma 4.7,

$$U_1^*(T) \ll xT^{\frac{1}{k}-1+\varepsilon},\ U_2^*(T) \ll T^{3-\frac{2}{k}}x^{\frac{2}{k}-1+\varepsilon} + x^{\frac{4}{k+1}-1+\varepsilon}T^2. \tag{4.29}$$

Hence, by (4.26), (4.27), (4.29) and Lemma 4.10,

$\int_{\mathfrak{M}} \mid S^*(\alpha)\Delta(\alpha) \mid d\alpha \ll \log x \max\limits_{1\le T\le x/R}(xT^{\frac{1}{k}-1+\varepsilon})^{1/2}(T^{3-\frac{2}{k}}x^{\frac{2}{k}-1+\varepsilon} + x^{\frac{4}{k+1}-1+\varepsilon}T^2)^{1/2} + x^{\frac{2}{k+1}+\varepsilon}$

$\ll x^{1+\frac{1}{2k}+\varepsilon}R^{\frac{1}{2k}-1} + x^{\frac{2}{k+1}+\frac{1}{2k}+\frac{1}{2}+\varepsilon}R^{-\frac{1}{2k}-\frac{1}{2}} + x^{\frac{2}{k+1}+\varepsilon}.$

Combining these with Lemma 4.5,

$\sum\limits_{q\le x/R}\sum\limits_{a=1}^{q}{}' \int_{\mathfrak{M}(q,a)} \mid S^*(\alpha;q,a)\Delta(\alpha;q,a)H_q(\alpha) \mid d\alpha$

$$\ll x^{\frac{3}{2}+\frac{1}{2k}+\varepsilon}R^{\frac{1}{2k}-1} + x^{1+\frac{2}{k+1}+\frac{1}{2k}+\varepsilon}R^{-\frac{1}{2k}-\frac{1}{2}} + x^{\frac{1}{2}+\frac{2}{k+1}+\varepsilon}. \tag{4.30}$$

LEMMA 4.11. We have

$$\sum_{q\le x/R}\sum_{a=1}^{q}{}' \int_{\mathfrak{M}(q,a)} \mid \Delta(\alpha;q,a) \mid^2 \mid K(\alpha) \mid d\alpha \ll x^{1+\frac{1}{k}-\frac{\tau}{2}} + x^{\frac{1}{2}+\frac{4}{k+1}-\frac{\tau}{2}}.$$

Proof. We use the method of (4.8) in [9]. Let $\mathfrak{M}_0(1,1) = \{\alpha : \mid \alpha - 1 \mid \le 1/x\}$ and for $j \ge 1$

$$\mathfrak{M}_j(q,a) = \begin{cases} \{\alpha : 2^{j-1}x^{-1} < \mid q\alpha - a \mid \le 2^j x^{-1}\}, & \text{when } \ q \le 2^{j-1}; \\ \{\alpha : \mid q\alpha - a \mid \le 2^j x^{-1}\}, & \text{when } \ 2^{j-1} < q \le 2^j. \end{cases}$$

$$\mathfrak{M}_j = \bigcup_{\substack{1\le a\le q\le 2^j \\ (a,q)=1}} \mathfrak{M}_j(q,a).$$

Choose J so that $2^{J-1} < x/R \le 2^J$. For $1 \le a \le q \le x/R,\ (a,q) = 1,$

then

$\mathfrak{M}(q,a) \subseteq \bigcup\limits_{\substack{0\le j\le J \\ q\le 2^j}} \mathfrak{M}_j(q,a), \quad \mathfrak{M} \subseteq \bigcup\limits_{0\le j\le J} \mathfrak{M}_j.$

By Lemme 4.4, when $\alpha \in \mathfrak{M}_j(q,a)$ with $1 \le a \le q \le 2^j, (a,q) = 1,$

$$K_q(\alpha) \ll 2^{-j}x \log x. \tag{4.31}$$

By Lemma 4.6 and (4.31),

$$\sum_{q\le x/R}\sum_{a=1}^{q}{}' \int_{\mathfrak{M}(q,a)} \mid \Delta(\alpha;q,a)\mid^2\mid K_q(\alpha)\mid d\alpha \ll x^{\frac{2}{k}+\varepsilon}(\tfrac{x}{R})^{2-\frac{2}{k}} + x^{\frac{4}{k+1}+\varepsilon}(\tfrac{x}{R})$$

$$\ll x^{2+\varepsilon}R^{\frac{2}{k}-2} + x^{1+\frac{4}{k+1}+\varepsilon}R^{-1}. \tag{4.32}$$

By Lemma 4.2, Lemma 4.9 and (4.32), the lemma follows.

By (4.23), (4.24) and Lemma 4.2, when $\alpha \in \mathfrak{M}(q,a)$,

$\mid S(\alpha)\mid^2 K(\alpha) =\mid S^*(\alpha;q,a)\mid^2 K(\alpha)$

$+\, 2(\Re(\overline{S^*(\alpha;q,a)}\Delta(\alpha;q,a)))(K_q(\alpha)+H_q(\alpha))+ \mid \Delta(\alpha;q,a)\mid^2 K(\alpha).$

By (4.13), (4.30) and Lemma 4.11,

$$S_A(x,Q) = S_{A1}(x,Q) + S_{A2}(x,Q) + O(x^{1+\frac{1}{k}-\frac{\tau}{2}} + x^{\frac{1}{2}+\frac{4}{k+1}-\frac{\tau}{2}}), \tag{4.33}$$

where

$$S_{A1}(x,Q) = \sum_{q\le x/R}\sum_{a=1}^{q}{}' \int_{\mathfrak{M}(q,a)} \mid S^*(\alpha;q,a)\mid^2 K(\alpha)d\alpha, \tag{4.34}$$

$$S_{A2}(x,Q) = 2\sum_{q\le x/R}\sum_{a=1}^{q}{}' \int_{\mathfrak{M}(q,a)} (\Re(\overline{S^*(\alpha;q,a)}\Delta(\alpha;q,a)))K_q(\alpha)d\alpha,$$

by (4.21) and (4.23)

$$S_{A2}(x,Q) = 2\varrho \sum_{q\le x/R} q^{-1}f(q)\sum_{a=1}^{q}{}' \int_{-1/qR}^{1/qR} (\Re(\overline{H(q,a)I(\beta)}\Delta(\frac{a}{q}+\beta;q,a)))K_q(\beta)d\beta. \tag{4.35}$$

5 The estimate of $S_{A2}(x,Q)$

LEMMA 5.1. We have

$$\sum_{b=1}^{q} h(q,b)\sum_{r\mid q}\mu(\frac{q}{r})g(r,b) = q^{-1}\sum_{b=1}^{q} h(q,b)\sum_{a=1}^{q}{}'G(q,a)e(\frac{-ab}{q}).$$

Proof. By (4.20) and $\sum\limits_{a=1}^{q}{}'e(\frac{an}{q}) = \sum\limits_{r\mid(q,n)} r\mu(\frac{q}{r})$,

$q^{-1}\sum\limits_{b=1}^{q} h(q,b)\sum\limits_{a=1}^{q}{}'G(q,a)e(\frac{-ab}{q})$

$= q^{-1}\sum\limits_{b=1}^{q} h(q,b)\sum\limits_{a=1}^{q}{}'e(\frac{-ab}{q})\sum\limits_{t=1}^{q} g(q,t)e(\frac{at}{q})$

$= q^{-1} \sum\limits_{b=1}^{q} h(q,b) \sum\limits_{r|q} r\mu(\frac{q}{r}) \sum\limits_{\substack{t=1 \\ r|t-b}}^{q} g(q,t).$

Hence

$$q^{-1}\sum_{b=1}^{q} h(q,b) \sum_{a=1}^{q}{}' G(q,a) e(\frac{-ab}{q}) = q^{-1}\sum_{r|q} r\mu(\frac{q}{r}) \sum_{b=1}^{q} h(q,b) \sum_{\substack{t=1 \\ r|t-b}}^{q} g(q,t). \tag{5.1}$$

By (4.14),

$$\sum_{\substack{t=1 \\ t\equiv b(r)}}^{q} g(q,t) = \sum_{n=1}^{\infty} \mu(n) \frac{(n^k,q)}{n^k} \sum_{\substack{t=1 \\ t\equiv b(r)}}^{q} \psi_k(n;q,t), \tag{5.2}$$

where

$$\psi_k(n;q,t) = \#\{u,v \geq 1 : uv = n,\ (u,v)=1,\ (u^k,q) \mid t,\ (v^k,q) \mid t+1\ \}.$$

Write $t = b + rl, q = rm$, and l runs through complete residues modulus m, we want to count the number of l.

We have

$(u^k, rm) \mid b + rl,\ (v^k, rm) \mid b + 1 + rl.$

Let

$$(u^k, r) = r_1,\ (v^k, r) = r_2, (r_1, r_2) = 1,$$

then

$r_1 \mid b,\ r_2 \mid b+1,$

$(\frac{u^k}{r_1}, m) \mid \frac{b}{r_1} + \frac{r}{r_1} l,\ (\frac{v^k}{r_2}, m) \mid \frac{b+1}{r_2} + \frac{r}{r_2} l.$

These congruences have a unique solution of l modulus $(\frac{u^k}{r_1}, m)(\frac{v^k}{r_2}, m)$, therefore the number of l is $\frac{m}{(\frac{u^k}{r_1},m)(\frac{v^k}{r_2},m)} = \frac{mr_1r_2}{(u^k,r_1m)(v^k,r_2m)}$.

Since $(r_1, r_2) = (r_1, v) = (r_2, u) = 1$, the number of l is

$\frac{mr_1r_2}{(u^k,r_1r_2m)(v^k,r_1r_2m)} = \frac{mr_1r_2}{(n^k,r_1r_2m)}$.

Hence

$\sum\limits_{\substack{t=1 \\ t\equiv b(r)}}^{q} g(q,t) = \sum\limits_{n=1}^{\infty} \mu(n) \frac{(n^k,q)}{n^k} \sum\limits_{u,v}{}^{*} \frac{mr_1r_2}{(n^k,r_1r_2m)},$

where $\sum\limits_{u,v}{}^{*}$ means $(u^k, r) = r_1 \mid b,\ (v^k, r) = r_2 \mid b{+}1,\ uv = n, (u, v) = 1,\ u, v \geq 1, q = rm.$

Let $r = r_1 r_2 \Delta$, then

$(q, n^k) = (r_1 r_2 \Delta m, u^k v^k) = (r_1 r_2 \Delta m, u^k)(r_1 r_2 \Delta m, v^k) = (r_1 \Delta m, u^k)(r_2 \Delta m, v^k)$

$= r_1 r_2 (\Delta m, \frac{u^k}{r_1})(\Delta m, \frac{v^k}{r_2}).$

Since $(\frac{u^k}{r_1}, r_2 \Delta) = (\frac{v^k}{r_2}, r_1 \Delta) = 1$, we have $(\Delta, \frac{u^k}{r_1} \frac{v^k}{r_2}) = 1$,

hence

$(q, n^k) = r_1 r_2 (m, \frac{u^k}{r_1})(m, \frac{v^k}{r_2}) = (r_1 r_2 m, u^k)(r_1 r_2 m, v^k) = (r_1 r_2 m, (uv)^k).$

So, by (5.1) and (5.2),

$q^{-1} \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{a=1}^{q}{}' G(q, a) e(\frac{-ab}{q}) = \sum\limits_{r|q} \mu(\frac{q}{r}) \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{n=1}^{\infty} \mu(n) \frac{1}{n^k} \sum\limits_{u,v}{}^{*} \frac{rmr_1 r_2}{q}$

$= \sum\limits_{r|q} \mu(\frac{q}{r}) \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{n=1}^{\infty} \mu(n) \frac{1}{n^k} \sum\limits_{u,v}{}^{*} r_1 r_2 = \sum\limits_{r|q} \mu(\frac{q}{r}) \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{n=1}^{\infty} \mu(n) \frac{(n^k, r)}{n^k} \psi_k(n; r, b),$

by (4.14), the lemma follows.

Write

$$L = \sum_{a=1}^{q}{}' \overline{H(q, a)} \Delta(\frac{a}{q} + \beta; q, a),$$

then, by (4.20) and (4.24),

$L = \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{a=1}^{q}{}' e(\frac{-ab}{q}) \Delta(\frac{a}{q} + \beta; q, a)$

$= \sum\limits_{n \leq x} (\mu_k(n) \mu_k(n{+}1) \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{a=1}^{q}{}' e(\frac{a}{q}(n - b)) - q^{-1} \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{a=1}^{q}{}' G(q, a) e(\frac{-ab}{q})) e(\beta n).$

Write

$U(y) = \sum\limits_{n \leq y} (\mu_k(n) \mu_k(n + 1) \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{a=1}^{q}{}' e(\frac{a}{q}(n - b)) - q^{-1} \sum\limits_{b=1}^{q} h(q, b) \sum\limits_{a=1}^{q}{}' G(q, a) e(\frac{-ab}{q}))$

$= U_1(y) - U_2(y).$

and using $e(n\beta) = e(x\beta) - 2\pi i \beta \int_n^x e(y\beta) dy$, we have

$$L = U(x) e(x\beta) - 2\pi i \beta \int_1^x U(y) e(y\beta) dy. \tag{5.3}$$

We estimate $U_1(y)$ first.

Write

$$E(x; q, a) = A_k(x; q, a) - q^{-1} g(q, a) x. \tag{5.4}$$

We have

$$U_1(y) = U_1([y]) = \sum_{b=1}^{q} h(q,b) \sum_{n\le[y]} \mu_k(n)\mu_k(n+1) \sum_{r|(q,n-b)} r\mu(\tfrac{q}{r})$$
$$= \sum_{r|q} r\mu(\tfrac{q}{r}) \sum_{b=1}^{q} h(q,b) \sum_{\substack{n\le[y]\\ r|n-b}} \mu_k(n)\mu_k(n+1)$$
$$= \sum_{r|q} \mu(\tfrac{q}{r}) \sum_{b=1}^{q} h(q,b)g(r,b)[y] + \sum_{r|q} r\mu(\tfrac{q}{r}) \sum_{b=1}^{q} h(q,b)E([y];r,b),$$

and

$$U_2(y) = \sum_{n\le[y]} q^{-1} \sum_{b=1}^{q} h(q,b) {\sum_{a=1}^{q}}' G(q,a)e(\tfrac{-ab}{q})$$
$$= q^{-1} \sum_{b=1}^{q} h(q,b) {\sum_{a=1}^{q}}' G(q,a)e(\tfrac{-ab}{q})[y].$$

We need an obvious result on $Y_k(x,Q)$.

LEMMA 5.2. For all Q, x, we have

$$Y_k(x,Q) \ll x^2 \log Q + Q^2.$$

Proof. By (1.4) and $g(q,a) \ll 1$,

$$Y_k(x,Q) \ll \sum_{q\le Q} \sum_{a=1}^{q} (x/q+1)^2 \ll \sum_{q\le Q} \sum_{a=1}^{q} (x^2q^{-2}+1) \ll x^2\log Q + Q^2,$$

the lemma follows.

By Lemma 5.1, we have

$$U(y) = \sum_{r|q} r\mu(\tfrac{q}{r}) \sum_{b=1}^{q} h(q,b)E([y];r,b).$$

Hence

$$L \ll \sum_{r|q} r\mu^2(\tfrac{q}{r}) \sum_{b=1}^{q} h(q,b) \mid E([x];r,b) \mid + \mid \beta \mid \int_1^x \sum_{r|q} r\mu^2(\tfrac{q}{r}) \sum_{b=1}^{q} h(q,b) \mid E([y];r,b) \mid dy.$$

By (4.35), Lemma 4.4 and $I(\beta) \ll x/(1+x \mid \beta \mid)$

$$S_{A2}(x,Q) \ll \sum_{q\le x/R} q^{-1} \int_0^{1/qR} \tfrac{x}{(1+x\beta)} \tfrac{x\log x}{q(1+x\beta)} (\sum_{r|q} r\mu^2(\tfrac{q}{r}) \sum_{b=1}^{q} h(q,b) \mid E([x];r,b) \mid$$
$$+ \beta \int_1^x \sum_{r|q} r\mu^2(\tfrac{q}{r}) \sum_{b=1}^{q} h(q,b) \mid E([y];r,b) \mid dy) d\beta \ll \int_0^{R^{-1}} \tfrac{x^2\log x}{(1+x\beta)^2} M(\beta) d\beta,$$

where

$$M(\beta) \ll$$

$$\sum_{\substack{q\le x/R\\ q\le 1/\beta R}} q^{-2} (\sum_{r|q} r\mu^2(\frac{q}{r}) \sum_{b=1}^{q} h(q,b) \mid E([x];r,b) \mid + \beta \int_1^x \sum_{r|q} r\mu^2(\frac{q}{r}) \sum_{b=1}^{q} h(q,b) \mid E([y];r,b) \mid dy). \tag{5.5}$$

When $r \mid q$, we have $\sum\limits_{b=1}^{q} \mid E(y;r,b) \mid = qr^{-1}\sum\limits_{b=1}^{r} \mid E(y;r,b) \mid$, then

$$\sum_{\substack{q\le x/R\\ q\le 1/\beta R}} q^{-2}\sum_{r|q} r\mu^2(\tfrac{q}{r})\sum_{b=1}^{q} h(q,b)\mid E([x];r,b)\mid \ll \sum_{r\le x/R}\sum_{b=1}^{r}\mid E([x];r,b)\mid \sum_{\substack{q\le x/R\\ r|q}} q^{-1}$$

$$\ll \log x \sum_{r\le x/R} r^{-1}\sum_{b=1}^{r}\mid E([x];r,b)\mid .$$

By Cauchy inequality and (1.5),

$$\sum_{\substack{q\le x/R\\ q\le 1/\beta R}} q^{-2}\sum_{r|q} r\mu^2(\tfrac{q}{r})\sum_{b=1}^{q} h(q,b)\mid E([x];r,b)\mid$$

$$\ll \log x(\sum_{r\le x/R}\sum_{b=1}^{r} r^{-2})^{1/2}(\sum_{r\le x/R}\sum_{b=1}^{r}\mid E([x];r,b)\mid^2)^{1/2}$$

$$\ll \log^{3/2} x(x^{\frac{2}{k}+\varepsilon}(\tfrac{x}{R})^{2-\frac{2}{k}}+x^{\frac{4}{k+1}+\varepsilon})^{1/2} \ll x^{1+\varepsilon}R^{\frac{1}{k}-1}+x^{\frac{2}{k+1}+\varepsilon},$$

this contributes to $S_{A2}(x,Q)$

$$\ll (x^{1+\varepsilon}R^{\frac{1}{k}-1}+x^{\frac{2}{k+1}+\varepsilon})\int_0^{R^{-1}}\tfrac{x^2\log x}{(1+x\beta)^2}d\beta$$

$$\ll x\log x(x^{1+\varepsilon}R^{\frac{1}{k}-1}+x^{\frac{2}{k+1}+\varepsilon}) \ll x^{2+\varepsilon}R^{\frac{1}{k}-1}+x^{1+\frac{2}{k+1}+\varepsilon}.$$

For the second term of $M(\beta)$, we have

$$\sum_{q\le x/R} q^{-2}\beta\int_1^x \sum_{r|q} r\mu^2(\tfrac{q}{r})\sum_{b=1}^{q} h(q,b)\mid E([y];r,b)\mid dy$$

$$\ll \beta\int_1^x \log x\sum_{r\le x/R} r^{-1}\sum_{b=1}^{r}\mid E([y];r,b)\mid dy$$

$$\ll \beta\int_1^{x/R}\log x\sum_{r\le x/R} r^{-1}\sum_{b=1}^{r}\mid E([y];r,b)\mid dy+\beta\int_{x/R}^{x}\log x\sum_{r\le x/R} r^{-1}\sum_{b=1}^{r}\mid E([y];r,b)\mid dy.$$

By Cauchy inequality and Lemma 5.2,

$$\beta\int_1^{x/R}\log^{3/2} x(\sum_{r\le x/R}\sum_{b=1}^{r}\mid E([y];r,b)\mid^2)^{1/2}dy$$

$$\ll \beta\int_1^{x/R}\log^{3/2} x(y^2\log x+(\tfrac{x}{R})^2)^{1/2}dy$$

$$\ll x^2R^{-2}\beta\log^2 x,$$

this contributes to $S_{A2}(x,Q)$

$$\ll x^2R^{-2}\log^2 x\int_0^{R^{-1}}\tfrac{x^2\log x}{(1+x\beta)^2}\beta d\beta \ll x^2R^{-2}\log^4 x.$$

By Cauchy inequality and (1.5),

$$\beta\int_{x/R}^{x}\log x\sum_{r\le x/R} r^{-1}\sum_{b=1}^{r}\mid E([y];r,b)\mid dy$$

$$\ll \beta\int_{x/R}^{x}\log^{3/2} x(x^{\frac{2}{k}+\varepsilon}(\tfrac{x}{R})^{2-\frac{2}{k}}+x^{\frac{4}{k+1}+\varepsilon})^{1/2}dy \ll \beta(x^{2+\varepsilon}R^{\frac{1}{k}-1}+x^{1+\frac{2}{k+1}+\varepsilon}),$$

this contributes to $S_{A2}(x,Q)$

$$\ll x^{2+\varepsilon}R^{\frac{1}{k}-1}+x^{1+\frac{2}{k+1}+\varepsilon}.$$

Consequently,

$$S_{A2}(x,Q)\ll x^{2+\varepsilon}R^{\frac{1}{k}-1}+x^{1+\frac{2}{k+1}+\varepsilon}+x^2R^{-2}\log^4 x. \tag{5.6}$$

6 The estimate of $S_{A1}(x,Q)$

By (4.2), (4.23) and (4.34),

$S_{A1}(x,Q)=\sum\limits_{q\le x/R}\sum\limits_{a=1}^{q}{}' \int_{-1/2}^{1/2} q^{-2}\mid G(q,a)I(\beta)\mid^2 K(\frac{a}{q}+\beta)d\beta -$

$\sum\limits_{q\le x/R}\sum\limits_{a=1}^{q}{}' \int_{1/qR\le|\beta|\le 1/2} q^{-2}\mid G(q,a)I(\beta)\mid^2 K(\frac{a}{q}+\beta)d\beta$

$$=S_B(x,Q)-S_C(x,Q), \tag{6.1}$$

say.

We consider $S_C(x,Q)$ first. By Lemma 4.4,

$S_C(x,Q)\ll\sum\limits_{q\le x/R}q^{-2}\sum\limits_{a=1}^{q}{}'\mid G(q,a)\mid^2\int_{1/qR\le|\beta|\le 1/2}\min(x^2,\beta^{-2})\sum\limits_{u\le\sqrt{x}}\frac{x}{u+x\|u\frac{a}{q}+u\beta\|}d\beta$

$\ll\sum\limits_{q\le x/R}q^{-2}\sum\limits_{a=1}^{q}{}'\mid G(q,a)\mid^2\sum\limits_{u\le\sqrt{x}}\int_{1/qR\le|\beta|\le 1/2}\frac{x\beta^{-2}}{u+x\|u\frac{a}{q}+u\beta\|}d\beta$

$\ll\sum\limits_{q\le x/R}q^{-2}\sum\limits_{a=1}^{q}{}'\mid G(q,a)\mid^2\sum\limits_{u\le\sqrt{x}}\int_{u/qR\le|\gamma|\le u/2}u\frac{x\gamma^{-2}}{u+x\|u\frac{a}{q}+\gamma\|}d\gamma.$

Now,

$ux\sum\limits_{0<|j|\le u/2}\int_{j-1/2\le\gamma\le j+1/2}\frac{\gamma^{-2}}{u+x\|u\frac{a}{q}+\gamma\|}d\gamma\ll ux\sum\limits_{0<|j|\le u/2}\int_{-1/2\le\gamma\le 1/2}\frac{j^{-2}}{u+x\|u\frac{a}{q}+\beta+j\|}d\beta$

$\ll ux\sum\limits_{0<|j|\le u/2}j^{-2}\int_{-1/2\le\gamma\le 1/2}\frac{1}{u+x|\beta|}d\beta\ll u\log x.$

Hence

$S_C(x,Q)\ll\sum\limits_{q\le x/R}q^{-2}\sum\limits_{a=1}^{q}{}'\mid G(q,a)\mid^2\sum\limits_{u\le\sqrt{x}}u\log x$

$+\sum\limits_{q\le x/R}q^{-2}\sum\limits_{a=1}^{q}{}'\mid G(q,a)\mid^2\sum\limits_{u\le\sqrt{x}}\int_{u/qR\le|\gamma|\le 1/2}u\frac{x\gamma^{-2}}{u+x\|u\frac{a}{q}+\gamma\|}d\gamma$

$=S_{C1}(x,Q)+S_{C2}(x,Q),$ say.

By Lemma 4.2 of [2], for any $Q\ge 1$,

$$\sum_{Q<q\le 2Q} q^{-2}f^2(q)H(q) \ll Q^{\frac{1}{k}-1+\varepsilon}. \tag{6.2}$$

Hence

$$S_{C1}(x,Q) \ll x\log x \sum_{q\le x/R} q^{-2}f^2(q)H(q) \ll x\log x. \tag{6.3}$$

LEMMA 6.1. We have

$$S_{C2}(x,Q) \ll x^{\frac{3}{2}+\frac{1}{2k}+2\tau}.$$

Proof. We have

$S_{C2}(x,Q) \ll \sum\limits_{q\le x/R} q^{-2} \sum\limits_{a=1}^{q}{}' \mid G(q,a)\mid^2 \sum\limits_{u\le\sqrt{x}} x\int_{u/qR\le|\gamma|\le 1/2}\gamma^{-2}d\gamma$

$\ll \sum\limits_{q\le x/R} q^{-2} \sum\limits_{a=1}^{q}{}' \mid G(q,a)\mid^2 \sum\limits_{u\le\sqrt{x}} x\frac{qR}{u}$

$\ll xR\log x \sum\limits_{q\le x/R} q^{-1} \sum\limits_{a=1}^{q}{}' \mid G(q,a)\mid^2 .$

By partial summation and (6.2),

$S_{C2}(x,Q) \ll xR\log x \sum\limits_{q\le x/R} q^{-1}H(q)f(q)^2 \ll xR(\frac{x}{R})^{\frac{1}{k}+\varepsilon} \ll x^{\frac{3}{2}+\frac{1}{2k}+2\tau},$

the lemma follows.

By Lemma 6.1 and (6.3),

$$S_C(x,Q) \ll x^{\frac{3}{2}+\frac{1}{2k}+2\tau}.$$

Hence

$$S_{A1}(x,Q) = S_B(x,Q) + O(x^{\frac{3}{2}+\frac{1}{2k}+2\tau}). \tag{6.4}$$

7 The estimate of $S_B(x,Q)$

We have

$\int_{-1/2}^{1/2} \mid I(\beta)\mid^2 K(\frac{a}{q}+\beta)d\beta = \sum\limits_{n_1,n_2\le x} \sum\limits_{\substack{uv\le x, u\le Q\\ n_1-n_2+uv=0}} e(uv\frac{a}{q}).$

By (6.1),

$S_B(x,Q) = \sum\limits_{q\le x/R} q^{-2} \sum\limits_{a=1}^{q}{}' \mid G(q,a)\mid^2 \sum\limits_{u\le Q}\sum\limits_{v\le x/u} e(uv\frac{a}{q}) \sum\limits_{\substack{m,n\le x\\ n=m+uv}} 1.$

Hence

$$S_B(x,Q)=\sum_{q\le x/R}q^{-2}\sum_{a=1}^{q}{}'\mid G(q,a)\mid^2\sum_{u\le Q}\sum_{v\le x/u}e(uv\frac{a}{q})[x-uv]. \qquad (7.1)$$

By (6.2),

$$\sum_{u\le Q}\sum_{v\le x/u}(x-uv)\sum_{q>x/R}q^{-2}\sum_{a=1}^{q}{}'\mid G(q,a)\mid^2$$
$$\ll\sum_{u\le Q}\sum_{v\le x/u}(x-uv)(x/R)^{\frac{1}{k}-1+\varepsilon}\ll(x/R)^{\frac{1}{k}-1+\varepsilon}\sum_{u\le Q}u\sum_{v\le x/u}(x/u-v)$$
$$\ll(x/R)^{\frac{1}{k}-1+\varepsilon}\sum_{u\le Q}u(x/u)^2\ll x^{\frac{3}{2}+\frac{1}{2k}+2\tau},$$

and

$$\sum_{q\le x/R}q^{-2}\sum_{a=1}^{q}{}'\mid G(q,a)\mid^2\sum_{u\le Q}\sum_{v\le x/u}1\ll x\log x.$$

Hence

$$S_B(x,Q)=\sum_{u\le Q}\sum_{v\le x/u}(x-uv)\mathfrak{S}(uv)+O(x^{\frac{3}{2}+\frac{1}{2k}+2\tau})=S_E(x,Q)+O(x^{\frac{3}{2}+\frac{1}{2k}+2\tau}), \qquad (7.2)$$

where

$$\mathfrak{S}(n)=\sum_{q=1}^{\infty}\sum_{a=1}^{q}{}'q^{-2}\mid G(q,a)\mid^2 e(\frac{an}{q}), \qquad (7.3)$$

$$S_E(x,Q)=\sum_{u\le Q}\sum_{v\le x/u}(x-uv)\mathfrak{S}(uv). \qquad (7.4)$$

8 The formula for $\mathfrak{S}(n)$

Write

$$J(q)=J(q;n)=\sum_{a=1}^{q}{}'\mid H(q,a)\mid^2 e(\frac{an}{q}), \qquad (8.1)$$

then, by (4.21) and (7.3),

$$\mathfrak{S}(n)=\varrho^2\sum_{q=1}^{\infty}q^{-2}f(q)^2J(q).$$

By (6.2), $\mathfrak{S}(n)$ converges absolutely.

LEMMA 8.1. For any natural numbers $q_1,q_2,\ (q_1,q_2)=1$, we have

$$J(q_1q_2) = J(q_1)J(q_2),$$

the function $J(q)$ is multiplicative.

Proof. By (4.19) and $h(q, a+mq) = h(q, a)$, write $a = a_1q_2 + a_2q_1,\ b = b_1q_2 + b_2q_1$,

$J(q_1q_2)$
$= \sum_{a_1=1}^{q_1}{}' \sum_{a_2=1}^{q_2}{}' \mid \sum_{b_1=1}^{q_1}{}' h(q_1, b_1q_2)e(\frac{a_1q_2b_1}{q_1}) \mid^2 e(\frac{a_1n}{q_1}) \mid \sum_{b_2=1}^{q_2}{}' h(q_2, b_2q_1)e(\frac{a_2q_1b_2}{q_2}) \mid^2 e(\frac{a_2n}{q_2})$
$= \sum_{a_1=1}^{q_1}{}' \mid \sum_{b_1=1}^{q_1}{}' h(q_1, b_1q_2)e(\frac{a_1q_2b_1}{q_1}) \mid^2 e(\frac{a_1n}{q_1}) \sum_{a_2=1}^{q_2}{}' \mid \sum_{b_2=1}^{q_2}{}' h(q_2, b_2q_1)e(\frac{a_2q_1b_2}{q_2}) \mid^2 e(\frac{a_2n}{q_2})$
$= J(q_1)J(q_2),$

the lemma follows.

We need to give a formula for $J(p^t)$, here p is a prime number and $t \geq 1$.

We have

$$J(p^t) = \sum_{a=1}^{p^t} \mid H(p^t, a) \mid^2 e(\frac{an}{p^t}) - \sum_{\substack{a=1 \\ p|a}}^{p^t} \mid H(p^t, a) \mid^2 e(\frac{an}{p^t}) = J_1(p^t) - J_2(p^t), \qquad (8.2)$$

where

$$J_1(p^t) = \sum_{a=1}^{p^t} \mid H(p^t, a) \mid^2 e(\frac{an}{p^t}),$$

$$J_2(p^t) = \sum_{\substack{a=1 \\ p|a}}^{p^t} \mid H(p^t, a) \mid^2 e(\frac{an}{p^t}).$$

LEMMA 8.2. When $t > k$

$$J(p^t) = 0.$$

Proof. When $t > k$, by (4.20),

$J_1(p^t) = \sum_{a=1}^{p^t} \mid \sum_{b=1}^{p^t} h(p^t, b)e(\frac{ab}{p^t}) \mid^2 e(\frac{an}{p^t})$

$= \sum_{b_1,b_2=1}^{p^t} h(p^t, b_1)h(p^t, b_2) \sum_{a=1}^{p^t} e(\frac{a(b_1-b_2)}{p^t})e(\frac{an}{p^t})$

$= p^t \sum_{\substack{b_1,b_2=1 \\ b_1-b_2+n\equiv 0(p^t)}}^{p^t} h(p^t, b_1)h(p^t, b_2)$

$= p^t \sum_{b=1}^{p^t} h(p^t, b)h(p^t, b+n).$

Also

$J_2(p^t) = \sum_{b_1,b_2=1}^{p^t} h(p^t, b_1)h(p^t, b_2) \sum_{\substack{a=1\\p|a}}^{p^t} e(\frac{a(b_1-b_2+n)}{p^t})$

$= \sum_{b_1,b_2=1}^{p^t} h(p^t, b_1)h(p^t, b_2) \sum_{a=1}^{p^{t-1}} e(\frac{a(b_1-b_2+n)}{p^{t-1}})$

$= p^{t-1} \sum_{\substack{b_1,b_2=1\\b_1-b_2+n\equiv 0(p^{t-1})}}^{p^t} h(p^t, b_1)h(p^t, b_2).$

By (3.9) of [2], for $t > k$, one has $h(p^t, a) = h(p^t, b)$ whenever $a \equiv b(mod\ p^k)$.Hence

$J_2(p^t) = p^{t-1} \sum_{b=1}^{p^t} h(p^t, b)h(p^t, b+n) \sum_{\substack{b_2=1\\b_2\equiv b+n(p^{t-1})}}^{p^t} = p^t \sum_{b=1}^{p^t} h(p^t, b)h(p^t, b+n) = J_1(p^t),$

therefore

$$J(p^t) = J_1(p^t) - J_2(p^t) = 0,$$

the lemma follows.

From now on, we suppose $1 \le t \le k$.

By (4.7) of [2],

$$h(p^t, a) = \begin{cases} 1 - p^{t-k}, & \text{if } p^t \mid a(a+1), \\ 1, & \text{otherwise.} \end{cases} \tag{8.3}$$

For fixed p^t, n, write

$$\Phi_1 = \Phi_1(p^t) = \#\{b:\ p^t \mid b(b+1),\ b = 1, 2, ..., p^t\},$$

$$\Phi_2 = \Phi_2(p^t) = \#\{b:\ p^t \mid b(b+1),\ p^t \mid (b+n)(b+n+1),\ b = 1, 2, ..., p^t\}, \tag{8.4}$$

$$\Phi_3 = \Phi_3(p^t) = \#\{b:\ p^t \dagger b(b+1),\ p^t \mid (b+n)(b+n+1),\ b = 1, 2, ..., p^t\},$$

$$\Phi_4 = \Phi_4(p^t) = \#\{b:\ p^t \dagger b(b+1),\ b = 1, 2, ..., p^t\},$$

$$\Phi_5 = \Phi_5(p^t) = \#\{b_1, b_2:\ p^t \mid b_1(b_1+1),\ p^t \mid b_2(b_2+1), b_1-b_2+n \equiv 0(p^{t-1}),\ b_1, b_2 = 1, 2, ..., p^t\}.$$

(8.5)

LEMMA 8.3. Suppose that $1 \leq t \leq k$. Then

$$J_1(p^t) = p^{2t} - 4p^{2t-k} + p^{3t-2k}\Phi_2.$$

Proof. As in the proof of Lemma 8.1 and Lemma 8.2, by (8.3),

$J_1(p^t) = p^t \sum_{b=1}^{p^t} h(p^t, b)h(p^t, b+n) = p^t(J_A + J_B),$

where

$J_A = \sum_{\substack{b=1 \\ p^t|b(b+1)}}^{p^t} (1 - p^{t-k})h(p^t, b+n), \ J_B = \sum_{\substack{b=1 \\ p^t \dagger b(b+1)}}^{p^t} h(p^t, b+n).$

Firstly, we deal with J_A.

We note that $\Phi_1 = 2, \ \Phi_3 = 2 - \Phi_2, \Phi_4 = p^t - 2.$

By (8.3),

$J_A = (1 - p^{t-k})^2\Phi_2 + (1 - p^{t-k}) \sum_{\substack{b=1 \\ p^t|b(b+1), p^t \dagger (b+n)(b+n+1)}}^{p^t} 1$

$= (1 - p^{t-k})^2\Phi_2 + (1 - p^{t-k})(2 - \Phi_2) = 2(1 - p^{t-k}) + (1 - p^{t-k})(-p^{t-k})\Phi_2,$

hence

$$J_A = 2(1 - p^{t-k}) - p^{t-k}(1 - p^{t-k})\Phi_2.$$

Now, we deal with J_B. By (8.3),

$J_B = (1 - p^{t-k})\Phi_3 + \Phi_4 - \Phi_3 = p^t - 2 - p^{t-k}\Phi_3 = p^t - 2 - p^{t-k}(2 - \Phi_2),$

hence

$$J_B = p^t - 2 - 2p^{t-k} + p^{t-k}\Phi_2.$$

Therefore

$J_1(p^t) = p^t(2(1-p^{t-k})-p^{t-k}(1-p^{t-k})\Phi_2+p^t-2-2p^{t-k}+p^{t-k}\Phi_2) = p^{2t}-4p^{2t-k}+p^{3t-2k}\Phi_2,$

the lemma follows.

LEMMA 8.4. We have

$$J_2(p) = p^2(1 - 2p^{-k})^2.$$

Proof. By (4.20) and (8.2),

$J_2(p) = (\sum_{b=1}^{p} h(p,b))^2,$

and

$\sum_{b=1}^{p} h(p,b) = \sum_{\substack{b=1 \\ p|b(b+1)}}^{p} (1-p^{1-k}) + \sum_{\substack{b=1 \\ p\dagger b(b+1)}}^{p} 1 = 2(1-p^{1-k}) + p - 2 = p - 2p^{1-k}.$

Hence

$$J_2(p) = p^2(1-2p^{-k})^2,$$

the lemma follows.

LEMMA 8.5. Suppose that $1 < t \le k$. Then

$$J_2(p^t) = p^{t-1}(p^{t+1} - 4p^{t-k+1} + p^{2t-2k}\Phi_5).$$

Proof. By (4.20) and (8.2),

$J_2(p^t) = p^{t-1}M,$

where

$$M = \sum_{\substack{b_1,b_2=1 \\ b_1-b_2+n\equiv 0(p^{t-1})}}^{p^t} h(p^t,b_1)h(p^t,b_2).$$

By (8.3), we obtain

$M = M_A + M_B,$

where

$M_A = \sum_{\substack{b_1=1 \\ p^t|b_1(b_1+1)}}^{p^t} \sum_{\substack{b_2=1 \\ b_1-b_2+n\equiv 0(p^{t-1})}}^{p^t} (1-p^{t-k})h(p^t,b_2),$

$M_B = \sum_{\substack{b_1=1 \\ p^t\dagger b_1(b_1+1)}}^{p^t} \sum_{\substack{b_2=1 \\ b_1-b_2+n\equiv 0(p^{t-1})}}^{p^t} h(p^t,b_2).$

Again, by (8.3) and (8.5),

$M_A = M_{A1} + M_{A2},$

where

$M_{A1} = (1-p^{t-k})^2\Phi_5,$

$M_{A2} = (1-p^{t-k})(\Phi_6 - \Phi_5),$

and

$$\Phi_6 = \#\{b_1, b_2 :\ p^t \mid b_1(b_1+1),\ b_1 - b_2 + n \equiv 0(p^{t-1}),\ b_1, b_2 = 1, 2, ..., p^t\} = 2p.$$

Hence

$$M_{A2} = (1 - p^{t-k})(2p - \Phi_5),$$

and

$M_A = (1 - p^{t-k})^2\Phi_5 + (1 - p^{t-k})(2p - \Phi_5) = 2p(1 - p^{t-k}) - p^{t-k}(1 - p^{t-k})\Phi_5.$

Write

$$\Phi_7 = \#\{b_1, b_2 :\ p^t \dagger b_1(b_1+1),\ b_1 - b_2 + n \equiv 0(p^{t-1}),\ b_1, b_2 = 1, 2, ..., p^t\},$$

$\Phi_8 = \#\{b_1, b_2 :\ p^t \dagger b_1(b_1+1),\ p^t \mid b_2(b_2+1),\ b_1 - b_2 + n \equiv 0(p^{t-1}),\ b_1, b_2 = 1, 2, ..., p^t\},$

then

$$\Phi_7 = p(p^t - 2), \quad \Phi_8 = 2p - \Phi_5.$$

By (8.3),

$M_B = M_{B1} + M_{B2},$

where

$M_{B1} = (1 - p^{t-k})\Phi_8, \quad M_{B2} = \Phi_7 - \Phi_8.$

Hence

$M_B = (1 - p^{t-k})\Phi_8 + \Phi_7 - \Phi_8 = p(p^t - 2) - p^{t-k}\Phi_8 = p(p^t - 2) - p^{t-k}(2p - \Phi_5).$

We obtain

$$M_B = p(p^t - 2) - 2p^{t-k+1} + p^{t-k}\Phi_5,$$

and

$M = 2p(1 - p^{t-k}) - p^{t-k}(1 - p^{t-k})\Phi_5 + p(p^t - 2) - 2p^{t-k+1} + p^{t-k}\Phi_5$

$= p^{t+1} - 4p^{t-k+1} + p^{2t-2k}\Phi_5,$

the lemma follows.

LEMMA 8.6 We have

$$\Phi_2=\begin{cases} 1, & \text{if } p^t \mid n(n+1), p^t \dagger n(n-1), \\ 1, & \text{if } p^t \dagger n(n+1), p^t \mid n(n-1), \\ 2, & \text{if } p^t \mid n(n+1), p^t \mid n(n-1), \\ 0, & \text{if } p^t \dagger n(n+1), p^t \dagger n(n-1). \end{cases}$$

Proof. By (8.4),

$$\Phi_2=\#\{b:\ p^t \mid b(b+1),\ p^t \mid (b+n)(b+n+1),\ b=1,2,...,p^t\},$$

the condition $p^t \mid b(b+1)$ means $b=p^t, p^t-1$. If $p^t \mid n(n+1)$,then $b=p^t$ satisfies the condition $p^t \mid (b+n)(b+n+1)$; If $p^t \mid n(n-1)$,then $b=p^t-1$ satisfies the condition $p^t \mid (b+n)(b+n+1)$; If $p^t \dagger n(n+1)$ and $p^t \dagger n(n-1)$, then neither $b=p^t$ nor $b=p^t-1$ satisfies the condition $p^t \mid (b+n)(b+n+1)$. The lemma follows.

LEMMA 8.7 We have

if $t=1$, then $\Phi_5=4$; if $1<t\leq k$, then

$$\Phi_5=\begin{cases} 2, & \text{if } p^{t-1} \mid n, \\ 2, & \text{if } p^{t-1} \dagger n, p^{t-1} \mid (n+1), p^{t-1} \mid (n-1), \\ 1, & \text{if } p^{t-1} \dagger n, p^{t-1} \mid (n+1), p^{t-1} \dagger (n-1), \\ 1, & \text{if } p^{t-1} \dagger n, p^{t-1} \dagger (n+1), p^{t-1} \mid (n-1), \\ 0, & \text{if } p^{t-1} \dagger n, p^{t-1} \dagger (n+1), p^{t-1} \dagger (n-1). \end{cases}$$

Proof. By (8.5),

$$\Phi_5=\#\{b_1,b_2:\ p^t \mid b_1(b_1+1),\ p^t \mid b_2(b_2+1), b_1-b_2+n\equiv 0(p^{t-1}),\ b_1,b_2=1,2,...,p^t\},$$

we need to count the number of b_1, b_2 that satisfy the conditions $(*)$: $p^t \mid b_1(b_1+1),\ p^t \mid b_2(b_2+1), b_1-b_2+n\equiv 0(p^{t-1})$.

We have $b_1, b_2 = p^t, p^t-1$.

If $t=1$, then $\Phi_5=4$ is valid. We suppose that $t>1$.

If $p^{t-1} \mid n$, then $b_1=b_2=p^t$ and $b_1=b_2=p^t-1$ satisfy the conditions $(*)$;

If $p^{t-1} \dagger n,\ p^{t-1} \mid (n+1), p^{t-1} \mid (n-1)$, then $b_1=p^t, b_2=p^t-1$ and $b_1=p^t-1, b_2=p^t$ satisfy the conditions $(*)$;

If $p^{t-1} \dagger n,\ p^{t-1} \mid (n+1), p^{t-1} \dagger (n-1)$, then only $b_1=p^t, b_2=p^t-1$ satisfies the conditions $(*)$; if $p^{t-1} \dagger n,\ p^{t-1} \dagger (n+1), p^{t-1} \mid (n-1)$ then only $b_1=p^t-1, b_2=p^t$

satisfies the conditions $(*)$;

if $p^{t-1} \dagger n,\ p^{t-1} \dagger (n+1), p^{t-1} \dagger (n-1)$ then there are not b_1, b_2 satisfy the conditions $(*)$.

The lemma follows.

LEMMA 8.8 Suppose that $1 \leq t \leq k$. Then

$$J(p^t) = p^{3t-2k-1}(p\Phi_2 - \Phi_5).$$

Proof. By (8.2), Lemma 8.3 and Lemma 8.4,

$J(p) = J_1(p) - J_2(p) = p^2 - 4p^{2-k} + p^{3-2k}\Phi_2 - p^2(1-2p^{-k})^2 = p^{2-2k}(p\Phi_2 - 4)$.

We suppose $1 < t \leq k$. By (8.2), Lemma 8.3 and Lemma 8.5,

$J(p^t) = J_1(p^t) - J_2(p^t) = p^{2t} - 4p^{2t-k} + p^{3t-2k}\Phi_2 - p^{t-1}(p^{t+1} - 4p^{t-k+1} + p^{2t-2k}\Phi_5)$

$= p^{3t-2k-1}(p\Phi_2 - \Phi_5)$,

the lemma follows.

By Lemma 8.1, Lemma 8.2 and Lemma 8.8,

$\mathfrak{S}(n) = \varrho^2 \prod\limits_p (1 + \sum\limits_{t=1}^{k} p^{-2t} f(p^t)^2 J(p^t)) = \varrho^2 \prod\limits_p (1 + \sum\limits_{t=1}^{k} p^{-2t} (\frac{p^k}{p^k-2})^2 p^{3t-2k-1}(p\Phi_2 - \Phi_5))$

$= \varrho^2 \prod\limits_p (1 + (p^k-2)^{-2} \sum\limits_{t=1}^{k} p^{t-1}(p\Phi_2 - \Phi_5))$,

that is

$$\mathfrak{S}(n) = \varrho^2 \prod_p \Delta_p, \tag{8.6}$$

where

$$\Delta_p = 1 + (p^k-2)^{-2} \sum_{t=1}^{k} p^{t-1}(p\Phi_2 - \Phi_5). \tag{8.7}$$

Write

$$\Pi_1 = \prod_{p|n(n+1)} \Delta_p, \quad \Pi_{1A} = \prod_{\substack{p|n(n+1) \\ p|n(n-1)}} \Delta_p, \quad \Pi_{1B} = \prod_{\substack{p|n(n+1) \\ p\dagger n(n-1)}} \Delta_p, \quad \Pi_1 = \Pi_{1A}\Pi_{1B}, \tag{8.8}$$

$$\Pi_2 = \prod_{p\dagger n(n+1)} \Delta_p, \quad \Pi_{2A} = \prod_{\substack{p\dagger n(n+1) \\ p|n-1}} \Delta_p, \quad \Pi_{2B} = \prod_{\substack{p\dagger n(n+1) \\ p\dagger n-1}} \Delta_p, \quad \Pi_2 = \Pi_{2A}\Pi_{2B}, \tag{8.9}$$

then

$$\mathfrak{S}(n) = \varrho^2 \Pi_{1A} \Pi_{1B} \Pi_{2A} \Pi_{2B}. \tag{8.10}$$

For fixed n, p, we write

$$p^{\beta_1} \parallel n,\ p^{\beta_2} \parallel n+1,\ p^{\beta_3} \parallel n-1.$$

We consider Π_{1A} first.

LEMMA 8.9 We have

$$\Pi_{1A} = \prod_{\substack{p|n \\ p^k \dagger n}} (1 - 4(p^k - 2)^{-2}) \prod_{p^k|n} (1 + 2(p^k - 2)^{-1}) \prod_{\substack{2^k|n+1 \\ or 2^k|n-1}} (1 + (2^k - 4)(2^k - 2)^{-2}) \times$$
$$\prod_{\substack{2^k \dagger n+1 \\ 2^k \dagger n-1, 2 \dagger n}} (1 - 4(2^k - 2)^{-2}).$$

Proof. We have

$$\Pi_{1A} = \prod_{\substack{p^k \dagger n \\ p|n}} \Delta_p \prod_{p^k|n} \Delta_p \prod_{2 \dagger n} \Delta_2. \tag{8.11}$$

If $p \mid n(n+1),\ p \mid n(n-1)$ and $p \dagger n$, then $p = 2, \beta_1 = 0,\ \min(\beta_2, \beta_3) = 1$.

Hence, by Lemma 8.6 and Lemma 8.7,

$\Phi_2(2) = 2, \Phi_5(2) = 4$, and then $2\Phi_2(2) - \Phi_5(2) = 0$.

When $1 < t \le k, 2 \dagger n$,

$$\Phi_2(2^t) = \begin{cases} 1, & \text{if } t \le \max(\beta_2, \beta_3), \\ 0, & \text{if } t > \max(\beta_2, \beta_3). \end{cases}$$

$$\Phi_5(2^t) = \begin{cases} 2, & \text{if } t = 2, \\ 1, & \text{if } 2 < t \le \max(\beta_2, \beta_3) + 1, \\ 0, & \text{if } t > \max(\beta_2, \beta_3) + 1. \end{cases}$$

Write $\beta = \max(\beta_2, \beta_3) \ge 2$. We have

$$2\Phi_2(2^t) - \Phi_5(2^t) = \begin{cases} 0, & \text{if } t = 2, \\ 1, & \text{if } 2 < t \le \beta, \\ -1, & \text{if } t = 1 + \beta, \\ 0, & \text{if } 1 + \beta < t \le k. \end{cases}$$

Hence

$$\Delta_2 = \begin{cases} 1 + (2^k - 2)^{-2}(\sum\limits_{3 \le t \le \beta} 2^{t-1} - 2^{\beta}), & \text{if } \beta < k, \\ 1 + (2^k - 2)^{-2} \sum\limits_{3 \le t \le k} 2^{t-1}, & \text{if } \beta \ge k. \end{cases}$$

Then

$$\Delta_2 = \begin{cases} 1 - 4(2^k - 2)^{-2}, & \text{if } \beta < k, \\ 1 + (2^k - 4)(2^k - 2)^{-2}, & \text{if } \beta \ge k. \end{cases} \tag{8.12}$$

We suppose that $p \mid n(n+1),\ p \mid n(n-1)$ and $p \mid n$, then $\beta_2 = \beta_3 = 0$.

By Lemma 8.6 and Lemma 8.7,

$$p\Phi_2(p) - \Phi_5(p) = 2p - 4,$$

$$\Phi_2(p^t) = \begin{cases} 2, & \text{if } p^t \mid n, \\ 0, & \text{if } p^t \dagger n, \end{cases}$$

and for $1 < t \le k$,

$$\Phi_5(p^t) = \begin{cases} 2, & \text{if } p^{t-1} \mid n, \\ 0, & \text{if } p^{t-1} \dagger n. \end{cases}$$

Hence

$$p\Phi_2(p^t) - \Phi_5(p^t) = \begin{cases} 2p - 2, & \text{if } t \le \beta_1 \\ -2, & \text{if } t = 1 + \beta_1, \\ 0, & \text{if } 1 + \beta_1 < t \le k. \end{cases}$$

Then

$$\Delta_p = \begin{cases} 1 + (p^k - 2)^{-2}(2p - 4) + (p^k - 2)^{-2}(\sum\limits_{2 \le t \le \beta_1} p^{t-1}(2p - 2) - 2p^{\beta_1}), & \text{if } \beta_1 < k, \\ 1 + (p^k - 2)^{-2}(2p - 4) + (p^k - 2)^{-2}(\sum\limits_{2 \le t \le k} p^{t-1}(2p - 2)), & \text{if } \beta_1 \ge k. \end{cases}$$

Consequently

$$\Delta_p = \begin{cases} 1 - 4(p^k - 2)^{-2}, & \text{if } 1 \le \beta_1 < k, \\ 1 + 2(p^k - 2)^{-1}, & \text{if } \beta_1 \ge k. \end{cases} \tag{8.13}$$

By (8.11), (8.12) and (8.13), the lemma follows.

Secondly, we consider Π_{1B}.

LEMMA 8.10 We have

$$\Pi_{1B} = \prod_{\substack{p^k \dagger n+1 \\ p \mid n+1, p \dagger n-1}} (1 - 4(p^k - 2)^{-2}) \prod_{\substack{p^k \mid n+1 \\ p \dagger n-1}} (1 + (p^k - 4)(p^k - 2)^{-2}).$$

Proof. We have $p \mid n(n+1), p \dagger n(n-1), \beta_1 = \beta_3 = 0, \beta_2 \geq 1$, and by Lemma 8.6 and Lemma 8.7,

$$p\Phi_2(p) - \Phi_5(p) = p - 4,$$

for $1 < t \leq k$,

$$\Phi_2(p^t) = \begin{cases} 1, & \text{if } p^t \mid n+1, \\ 0, & \text{if } p^t \dagger n+1, \end{cases}$$

$$\Phi_5(p^t) = \begin{cases} 1, & \text{if } p^{t-1} \mid n+1, \\ 0, & \text{if } p^{t-1} \dagger n+1, \end{cases}$$

then

$$p\Phi_2(p^t) - \Phi_5(p^t) = \begin{cases} p-1, & \text{if } t \leq \beta_2 \\ -1, & \text{if } t = 1 + \beta_2, \\ 0, & \text{if } 1 + \beta_2 < t \leq k, \end{cases}$$

and

$$\Delta_p = \begin{cases} 1 + (p^k - 2)^{-2}(p - 4) + (p^k - 2)^{-2}(\sum\limits_{2 \leq t \leq \beta_2} p^{t-1}(p-1) - p^{\beta_2}), & \text{if } \beta_2 < k, \\ 1 + (p^k - 2)^{-2}(p - 4) + (p^k - 2)^{-2}(\sum\limits_{2 \leq t \leq k} p^{t-1}(p-1)), & \text{if } \beta_2 \geq k. \end{cases}$$

Consequently

$$\Delta_p = \begin{cases} 1 - 4(p^k - 2)^{-2}, & \text{if } 1 \leq \beta_2 < k, \\ 1 + (p^k - 4)(p^k - 2)^{-2}, & \text{if } \beta_2 \geq k. \end{cases} \tag{8.14}$$

By (8.8) and (8.14), the lemma follows.

Now, we consider Π_2. By $p \dagger n(n+1)$, Lemma 8.6 and Lemma 8.7,

$$\Phi_2(p^t) = \begin{cases} 1, & \text{if } p^t \mid n-1, \\ 0, & \text{if } p^t \dagger n-1, \end{cases} \tag{8.15}$$

and for $1 < t \leq k$

$$\Phi_5(p^t) = \begin{cases} 1, & \text{if } p^{t-1} \mid n-1, \\ 0, & \text{if } p^{t-1} \dagger n-1. \end{cases} \tag{8.16}$$

LEMMA 8.11 We have

$$\Pi_{2A} = \prod_{\substack{p\dagger n(n+1)\\ p^k\dagger n-1, p|n-1}} (1-4(p^k-2)^{-2}) \prod_{\substack{p\dagger n(n+1)\\ p^k|n-1}} (1+(p^k-4)(p^k-2)^{-2}).$$

Proof. We have $p \dagger n(n+1), p \mid (n-1), \beta_1 = \beta_2 = 0, \beta_3 \geq 1$, and by Lemma 8.6, Lemma 8.7, (8.15) and (8.16),

$$p\Phi_2(p) - \Phi_5(p) = p - 4,$$

and for $1 < t \leq k$, by (8.15) and (8.16),

$$p\Phi_2(p^t) - \Phi_5(p^t) = \begin{cases} p-1, & \text{if } t \leq \beta_3 \\ -1, & \text{if } t = 1+\beta_3, \\ 0, & \text{if } 1+\beta_3 < t \leq k, \end{cases}$$

and

$$\Delta_p = \begin{cases} 1+(p^k-2)^{-2}(p-4)+(p^k-2)^{-2}(\sum\limits_{2\leq t\leq \beta_3} p^{t-1}(p-1)-p^{\beta_3}), & \text{if } \beta_3 < k, \\ 1+(p^k-2)^{-2}(p-4)+(p^k-2)^{-2}(\sum\limits_{2\leq t\leq k} p^{t-1}(p-1)), & \text{if } \beta_3 \geq k. \end{cases}$$

Consequently

$$\Delta_p = \begin{cases} 1-4(p^k-2)^{-2}, & \text{if } 1 \leq \beta_3 < k, \\ 1+(p^k-4)(p^k-2)^{-2}, & \text{if } \beta_3 \geq k. \end{cases} \tag{8.17}$$

By (8.9) and (8.17), the lemma follows.

LEMMA 8.12 We have

$$\Pi_{2B} = \prod_{p\dagger n(n+1)(n-1)} (1-4(p^k-2)^{-2}).$$

Proof. We have $p \dagger n(n+1)(n-1), \beta_1 = \beta_2 = \beta_3 = 0$, and by Lemma 8.6, Lemma 8.7, (8.15) and (8.16),

$$p\Phi_2(p) - \Phi_5(p) = -4,$$

and for $1 < t \leq k$,

$$\Phi_2(p^t) = 0,\ \Phi_5(p^t) = 0,$$

and

$p\Phi_2(p^t) - \Phi_5(p^t) = 0$, and $\Delta_p = 1-4(p^k-2)^{-2}$,

the lemma follows.

LEMMA 8.13 We have

$$\mathfrak{S}(n) = \varrho^2 \prod_{\substack{p|n\\ p^k\dagger n}} (1-4(p^k-2)^{-2}) \prod_{p^k|n} (1+2(p^k-2)^{-1}) \prod_{\substack{2^k|n+1\\ or 2^k|n-1}} (1+(2^k-4)(2^k-2)^{-2}) \times$$
$$\prod_{\substack{2^k\dagger n+1\\ 2^k\dagger n-1, 2\dagger n}} (1-4(2^k-2)^{-2}) \prod_{\substack{p^k\dagger n+1\\ p|n+1, p\dagger n-1}} (1-4(p^k-2)^{-2}) \prod_{\substack{p^k|n+1\\ p\dagger n-1}} (1+(p^k-4)(p^k-2)^{-2}) \times$$
$$\prod_{\substack{p\dagger n(n+1)\\ p^k\dagger n-1, p|n-1}} (1-4(p^k-2)^{-2}) \prod_{\substack{p\dagger n(n+1)\\ p^k|n-1}} (1+(p^k-4)(p^k-2)^{-2}) \prod_{p\dagger n(n+1)(n-1)} (1-4(p^k-2)^{-2}).$$

Proof. By Lemma 8.9, Lemma 8.10, Lemma 8.11, Lemma 8.12 and (8.10), the lemma follows.

LEMMA 8.14 We have

$$\mathfrak{S}(n) = \varrho^2 \prod_{p\neq 2} (1-4(p^k-2)^{-2}) \prod_{\substack{2^k|n+1\\ or 2^k|n-1}} (1+(2^k-4)(2^k-2)^{-2}) \prod_{\substack{2^k\dagger n+1\\ 2^k\dagger n-1, 2\dagger n}} (1-4(2^k-2)^{-2}) \times$$
$$\prod_{\substack{2|n\\ 2^k\dagger n}} (1-4(2^k-2)^{-2}) \prod_{2^k|n} (1+2(2^k-2)^{-1}) \prod_{\substack{p^k|n\\ p\neq 2}} \frac{p^k-2}{p^k-4} \prod_{\substack{p^k|n+1\\ p\neq 2}} \frac{p^k-3}{p^k-4} \prod_{\substack{p^k|n-1\\ p\neq 2}} \frac{p^k-3}{p^k-4}.$$

Proof. By Lemma 8.13 and

$$\prod_{p\dagger n(n+1)(n-1)} (1-4(p^k-2)^{-2}) = \prod_{p\neq 2} (1-4(p^k-2)^{-2}) \prod_{\substack{p|n(n+1)(n-1)\\ p\neq 2}} (1-4(p^k-2)^{-2})^{-1},$$

the lemma follows at once.

9 The function $G(u;s)$

By (7.4),

$$S_E(x,Q) = \frac{1}{2\pi i} \int_{(2)} \sum_{u\leq Q} \sum_{v=1}^{\infty} \mathfrak{S}(uv)(uv)^{-s} \frac{x^{s+1}}{s(s+1)} ds, \tag{9.1}$$

where $\int_{(2)}$ means $\int_{2-i\infty}^{2+i\infty}$.

Write

$$G(u;s) = \sum_{v=1}^{\infty} \mathfrak{S}(uv) v^{-s}, \quad \Re s > 1,$$

then

$$S_E(x,Q) = \sum_{u\leq Q} \frac{1}{2\pi i} \int_{(2)} u^{-s} G(u;s) \frac{x^{s+1}}{s(s+1)} ds. \tag{9.2}$$

Let

$$G_1(u,s) = \sum_{\substack{v=1\\ 2^k|uv}}^{\infty} \mathfrak{S}(uv) v^{-s}, \; G_2(u,s) = \sum_{\substack{v=1\\ 2|uv, 2^k\dagger uv}}^{\infty} \mathfrak{S}(uv) v^{-s},$$

$$G_3(u,s) = \sum_{\substack{2\dagger uv, v=1\\ 2^k\dagger uv+1, 2^k\dagger uv-1}}^{\infty} \mathfrak{S}(uv)v^{-s},\ G_4(u,s) = \sum_{\substack{2\dagger uv, v=1\\ 2^k|uv+1 or 2^k|uv-1}}^{\infty} \mathfrak{S}(uv)v^{-s},$$

then

$$G(u;s) = G_1(u,s) + G_2(u,s) + G_3(u,s) + G_4(u,s). \tag{9.3}$$

As in [3], for integer a, let

$\xi_k = e^{2\pi i/k} = e(\frac{1}{k}),\ \zeta(s;\xi_k^a) = \sum_{m=1}^{\infty} \xi_k^{am} m^{-s},\ \zeta(s;a,k) = \sum_{n\equiv a(k)} n^{-s}, \Re s > 1,$

and for $0 < a \leq 1$, let $\zeta(s,a) = \sum_{n=0}^{\infty} (n+a)^{-s}, \Re s > 1$.

When $1 \leq a \leq k$, we have $\zeta(s;a,k) = k^{-s}\zeta(s, \frac{a}{k})$.

For odd integer, we define the multiplicative functions $g_1(d), g_2(d)$ as follows

$$g_1(p^t) = \frac{2}{p^k - 4},\quad g_2(p^t) = \frac{1}{p^k - 4},\ t \geq 1, p > 2, \tag{9.4}$$

then

$$\mu(d)^2 g_1(d) \ll \tau(d) d^{-k},\quad \mu(d)^2 g_2(d) \ll d^{-k}. \tag{9.5}$$

For fixed $u,\ d_1, d_2,\ d_3$ and $\Re s > 1$, write

$$H_1(s) = H_1(s;u,d_1,d_2,d_3) = \sum_{\substack{2^k|uv, d_1^k|uv\\ d_2^k|uv+1, d_3^k|uv-1}} v^{-s},$$

$$H_2(s) = H_2(s;u,d_1,d_2,d_3) = \sum_{\substack{2|uv, d_1^k|uv\\ d_2^k|uv+1, d_3^k|uv-1}} v^{-s},$$

$$H_3(s) = H_3(s;u,d_1,d_2,d_3) = \sum_{\substack{d_1^k|uv\\ d_2^k|uv+1, d_3^k|uv-1}} v^{-s},$$

$$H_4(s) = H_4(s;u,d_1,d_2,d_3) = \sum_{\substack{2^k|uv+1, d_1^k|uv\\ d_2^k|uv+1, d_3^k|uv-1}} v^{-s},$$

$$H_5(s) = H_5(s;u,d_1,d_2,d_3) = \sum_{\substack{2^k|uv-1, d_1^k|uv\\ d_2^k|uv+1, d_3^k|uv-1}} v^{-s}.$$

By Lemma 8.14 and (9.4),

$$\mathfrak{S}(n) = \varrho^2 \prod_{p\neq 2}(1 - 4(p^k-2)^{-2}) \prod_{\substack{2^k|n+1\\ or 2^k|n-1}} (1 + (2^k-4)(2^k-2)^{-2}) \times$$

$$\prod_{\substack{2^k\dagger n+1\\ 2^k\dagger n-1, 2\dagger n}} (1 - 4(2^k-2)^{-2}) \prod_{\substack{2|n\\ 2^k\dagger n}} (1 - 4(2^k-2)^{-2}) \prod_{2^k|n} (1 + 2(2^k-2)^{-1}) \times$$

$$\sum_{\substack{d_1^k|n\\(d_1,2)=1}} \mu(d_1)^2 g_1(d_1) \sum_{\substack{d_2^k|n+1\\(d_2,2)=1}} \mu(d_2)^2 g_2(d_2) \sum_{\substack{d_3^k|n-1\\(d_3,2)=1}} \mu(d_3)^2 g_2(d_3).$$

If we write

$$h(d_1, d_2, d_3) = \mu(d_1 d_2 d_3)^2 g_1(d_1) g_2(d_2) g_2(d_3),$$

then, by (9.5),

$$h(d_1, d_2, d_3) \ll \tau(d_1) d_1^{-k} d_2^{-k} d_3^{-k}, \tag{9.6}$$

and

$$\begin{aligned}\mathfrak{S}(n) = \varrho^2 \prod_{p\neq 2}(1-4(p^k-2)^{-2}) \prod_{\substack{2^k|n+1\\ or 2^k|n-1}} (1+(2^k-4)(2^k-2)^{-2})\times \\ \prod_{\substack{2^k\dagger n+1\\ 2^k\dagger n-1, 2\dagger n}} (1-4(2^k-2)^{-2}) \prod_{\substack{2|n\\ 2^k\dagger n}} (1-4(2^k-2)^{-2}) \prod_{2^k|n} (1+2(2^k-2)^{-1})\times \\ \sum_{\substack{(d_1 d_2 d_3,2)=1\\ d_1^k|n, d_2^k|n+1, d_3^k|n-1}} h(d_1, d_2, d_3).\end{aligned} \tag{9.7}$$

LEMMA 9.1 For $\Re s > 1$, we have

$G_1(u,s) = \varrho^2 \prod_{p\neq 2}(1-4(p^k-2)^{-2})(1+2(2^k-2)^{-1}) \sum_{(d_1 d_2 d_3,2)=1} h(d_1,d_2,d_3) H_1(s),$
$G_2(u,s) = \varrho^2 \prod_{p\neq 2}(1-4(p^k-2)^{-2})(1-4(2^k-2)^{-2}) \sum_{(d_1 d_2 d_3,2)=1} h(d_1,d_2,d_3)(H_2(s)-H_1(s)),$
$G_3(u,s) = \varrho^2 \prod_{p\neq 2}(1-4(p^k-2)^{-2})(1-4(2^k-2)^{-2}) \sum_{(d_1 d_2 d_3,2)=1} h(d_1,d_2,d_3)(H_3(s)-H_2(s)-$
$H_4(s)-H_5(s)),$

$G_4(u,s) = \varrho^2 \prod_{p\neq 2}(1-4(p^k-2)^{-2})(1+(2^k-4)(2^k-2)^{-2}) \sum_{(d_1 d_2 d_3,2)=1} h(d_1,d_2,d_3)(H_4(s)+$
$H_5(s)),$

$G(u;s) = \varrho^2 \prod_{p\neq 2}(1-4(p^k-2)^{-2}) \sum_{(d_1 d_2 d_3,2)=1} h(d_1,d_2,d_3)(2^{k+1}(2^k-2)^{-2} H_1(s) +$
$(1-4(2^k-2)^{-2}) H_3(s) + 2^k(2^k-2)^{-2} H_4(s) + 2^k(2^k-2)^{-2} H_5(s)).$

Proof. By (9.3) and (9.7), the lemma follows.

We only consider the case: $\mu(d_1 d_2 d_3)^2 = 1,\ (d_2 d_3, u) = 1.$ Write

$$\frac{2^k d_1^k u}{(u, 2^k d_1^k)} b_1 \equiv 1 (mod\ d_2^k),\ \frac{2^k d_1^k u}{(u, 2^k d_1^k)} b_2 \equiv 1 (mod\ d_3^k), \tag{9.8}$$

$$\begin{cases} d_3^k\overline{d_3^k} \equiv 1(mod\ d_2^k), \\ d_2^k\overline{d_2^k} \equiv 1(mod\ d_3^k), \end{cases} \tag{9.9}$$

$$A_1 \equiv -b_1 d_3^k\overline{d_3^k} + b_2 d_2^k\overline{d_2^k}\ (mod\ d_2^k d_3^k),\ 1 \le A_1 \le d_2^k d_3^k, (A_1, d_2 d_3) = 1, \tag{9.10}$$

$$\frac{d_1^k u}{(u, d_1^k)} b_5 \equiv 1(mod\ d_2^k),\ \frac{d_1^k u}{(u, d_1^k)} b_6 \equiv 1(mod\ d_3^k), \tag{9.11}$$

$$A_3 \equiv -b_5 d_3^k\overline{d_3^k} + b_6 d_2^k\overline{d_2^k}\ (mod\ d_2^k d_3^k),\ 1 \le A_3 \le d_2^k d_3^k, (A_3, d_2 d_3) = 1, \tag{9.12}$$

$$\frac{d_1^k u}{(u, d_1^k)} b_7 \equiv 1(mod\ 2^k d_2^k),\ \frac{d_1^k u}{(u, d_1^k)} b_8 \equiv 1(mod\ d_3^k), \tag{9.13}$$

$$\begin{cases} d_3^k\overline{(d_3^k)} \equiv 1(mod\ 2^k d_2^k), \\ 2^k d_2^k\overline{(2^k d_2^k)} \equiv 1(mod\ d_3^k), \end{cases} \tag{9.14}$$

$$A_4 \equiv -b_7 d_3^k\overline{(d_3^k)} + b_8 2^k d_2^k\overline{(2^k d_2^k)}\ (mod\ 2^k d_2^k d_3^k),\ 1 \le A_4 \le 2^k d_2^k d_3^k, (A_4, 2d_2 d_3) = 1, \tag{9.15}$$

$$\frac{d_1^k u}{(u, d_1^k)} b_9 \equiv 1(mod\ d_2^k),\ \frac{d_1^k u}{(u, d_1^k)} b_{10} \equiv 1(mod\ 2^k d_3^k), \tag{9.16}$$

$$\begin{cases} 2^k d_3^k\overline{\{2^k d_3^k\}} \equiv 1(mod\ d_2^k), \\ d_2^k\overline{\{d_2^k\}} \equiv 1(mod\ 2^k d_3^k), \end{cases} \tag{9.17}$$

$$A_5 \equiv -b_9 2^k d_3^k\overline{\{2^k d_3^k\}} + b_{10} d_2^k\overline{\{d_2^k\}}\ (mod\ 2^k d_2^k d_3^k),\ 1 \le A_5 \le 2^k d_2^k d_3^k, (A_5, 2d_2 d_3) = 1. \tag{9.18}$$

LEMMA 9.2 For $\Re s > 1$, we have

$H_1(s) = \frac{(u, 2^k d_1^k)^s}{2^{ks} d_1^{ks}} \zeta(s; A_1, d_2^k d_3^k),\ H_3(s) = \frac{(u, d_1^k)^s}{d_1^{ks}} \zeta(s; A_3, d_2^k d_3^k),$

$H_4(s) = \frac{(u,d_1^k)^s}{d_1^{ks}}\zeta(s; A_4, 2^k d_2^k d_3^k),\ H_5(s) = \frac{(u,d_1^k)^s}{d_1^{ks}}\zeta(s; A_5, 2^k d_2^k d_3^k).$

Proof. We have

$2^k \mid uv, d_1^k \mid uv, d_2^k \mid uv+1, d_3^k \mid uv-1.$

Write $a = \frac{2^k d_1^k}{(u,2^k d_1^k)}, v = av_1$, then

$$uav_1 \equiv -1(mod\ d_2^k),\ uav_1 \equiv 1(mod\ d_3^k).$$

By (9.8) and (9.9),

$$\begin{cases} v_1 \equiv -b_1(mod\ d_2^k), \\ v_1 \equiv b_2(mod\ d_3^k), \end{cases}$$

hence, by (9.10), we obtain $v_1 \equiv A_1(mod\ d_2^k d_3^k)$ and

$H_1(s) = a^{-s} \sum\limits_{v_1 \equiv A_1(d_2^k d_3^k)} v_1^{-s} = a^{-s}\zeta(s; A_1, d_2^k d_3^k).$

Similarly, the formulas for $H_3(s), H_4(s)$ and $H_5(s)$ follow from (9.11),(9.12),(9.13),(9.14), (9.15),(9.16),(9.17) and (9.18), and the lemma follows.

We use the following results of [3].

Let $G(s) = -i(2\pi)^{s-1}\Gamma(1-s)$, then

$$\zeta(s; a, k) = G(s)k^{-s}(e^{\frac{1}{2}i\pi s}\zeta(1-s; \xi_k^a) - e^{\frac{-1}{2}i\pi s}\zeta(1-s; \xi_k^{-a})). \tag{9.19}$$

By Lemma 9.1 and Lemma 9.2($\Re s > 1$),

$$\begin{aligned} G(u; s) = \varrho^2 \prod_{p \neq 2}(1 - 4(p^k-2)^{-2}) \sum_{\substack{(d_2 d_3, u)=1 \\ (d_1 d_2 d_3, 2)=1}} & h(d_1, d_2, d_3)\times \\ \{2^{k+1}(2^k-2)^{-2}\frac{(u,2^k d_1^k)^s}{2^{ks}d_1^{ks}}&\zeta(s; A_1, d_2^k d_3^k) \\ +(1-4(2^k-2)^{-2})\frac{(u,d_1^k)^s}{d_1^{ks}}&\zeta(s; A_3, d_2^k d_3^k) \\ +2^k(2^k-2)^{-2}\frac{(u,d_1^k)^s}{d_1^{ks}}&\zeta(s; A_4, 2^k d_2^k d_3^k) \\ +2^k(2^k-2)^{-2}\frac{(u,d_1^k)^s}{d_1^{ks}}&\zeta(s; A_5, 2^k d_2^k d_3^k)\}. \end{aligned} \tag{9.20}$$

We take (9.20) as the analytic continuation of $G(u; s), \Re s > \frac{1}{k} - 1$, and $G(u; s)$ has at most one simple pole at $s = 1$.

Write

$G_{(1)}(u, s) = \sum\limits_{\substack{(d_2 d_3, u)=1 \\ (d_1 d_2 d_3, 2)=1}} h(d_1, d_2, d_3)\frac{(u,2^k d_1^k)^s}{2^{ks}d_1^{ks}}\zeta(s; A_1, d_2^k d_3^k),$

$G_{(3)}(u,s) = \sum_{\substack{(d_2d_3,u)=1\\(d_1d_2d_3,2)=1}} h(d_1,d_2,d_3)\frac{(u,d_1^k)^s}{d_1^{ks}}\zeta(s;A_3,d_2^kd_3^k),$

$G_{(4)}(u,s) = \sum_{\substack{(d_2d_3,u)=1\\(d_1d_2d_3,2)=1}} h(d_1,d_2,d_3)\frac{(u,d_1^k)^s}{d_1^{ks}}\zeta(s;A_4,2^kd_2^kd_3^k),$

$G_{(5)}(u,s) = \sum_{\substack{(d_2d_3,u)=1\\(d_1d_2d_3,2)=1}} h(d_1,d_2,d_3)\frac{(u,d_1^k)^s}{d_1^{ks}}\zeta(s;A_5,2^kd_2^kd_3^k),$

then

$$\begin{aligned} G(u;s) &= \varrho^2\prod_{p\neq 2}(1-4(p^k-2)^{-2})2^{k+1}(2^k-2)^{-2}G_{(1)}(u,s)\\ &+\varrho^2\prod_{p\neq 2}(1-4(p^k-2)^{-2})(1-4(2^k-2)^{-2})G_{(3)}(u,s)\\ &+\varrho^2\prod_{p\neq 2}(1-4(p^k-2)^{-2})2^k(2^k-2)^{-2}G_{(4)}(u,s)\\ &+\varrho^2\prod_{p\neq 2}(1-4(p^k-2)^{-2})2^k(2^k-2)^{-2}G_{(5)}(u,s). \end{aligned} \tag{9.21}$$

10 The formula for $S_E(x,Q)$

Let $T>1$, by (9.2),

$S_E(x,Q) = \sum_{u\le Q}\frac{1}{2\pi i}(\int_{2+iT}^{2+i\infty}+\int_{2-iT}^{2+iT}+\int_{2-i\infty}^{2-iT})u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}ds.$

We have

$\sum_{u\le Q}\frac{1}{2\pi i}(\int_{2+iT}^{2+i\infty}+\int_{2-i\infty}^{2-iT})u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}ds \ll \sum_{u\le Q}u^{-2}x^3T^{-1} \ll x^3T^{-1},$

therefore

$$S_E(x,Q) = \sum_{u\le Q}\frac{1}{2\pi i}\int_{2-iT}^{2+iT}u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}ds + O(x^3T^{-1}) = A + O(x^3T^{-1}), \tag{10.1}$$

where

$A = \sum_{u\le Q}\frac{1}{2\pi i}\int_{2-iT}^{2+iT}u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}ds.$

Let $\frac{1}{k}-1+\tau\le\sigma<0$, by residue theorem

$A = \sum_{u\le Q}\frac{1}{2\pi i}\int_{\sigma-iT}^{\sigma+iT}u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}ds + \sum_{u\le Q}\frac{1}{2\pi i}\int_{2-iT}^{\sigma-iT}u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}ds$
$+\sum_{u\le Q}\frac{1}{2\pi i}\int_{\sigma+iT}^{2+iT}u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}ds + Res_{s=0,1}(\sum_{u\le Q}u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}).$

LEMMA 10.1 For fixed $\frac{1}{k}-1+\tau\le\sigma\le 2, \mid t\mid\ge 1$ and $s=\sigma+it$, we have the Stirling formula

$$\Gamma(s) = \sqrt{2\pi}e^{-\frac{\pi}{2}|t|}\mid t\mid^{\sigma-\frac{1}{2}}e^{it(\log|t|-1)}e^{\lambda\frac{i\pi}{2}(\sigma-\frac{1}{2})}(1+O(\mid t\mid^{-1})),$$

where $\lambda = 1$ if $t > 0$, and $\lambda = -1$ if $t < 0$; and

$$G(u;s) \ll | t | + | t |^{1-\sigma}, \qquad \frac{1}{k} - 1 + \tau \le \Re s \le 2, \ | t | \ge 1.$$

Proof. The first part is the Corollary 3.3.3 of [5], or see 4.42 of [6].

By Theorem 7.1.1 of [5], for $-1 < \Re s \le 2, \ y \ge 0$

$\zeta(s,a) = \sum\limits_{0 \le n \le y} \frac{1}{(n+a)^s} + \frac{(y+a)^{1-s}}{s-1} + \frac{y-[y]-\frac{1}{2}}{(y+a)^s} - s\int_y^\infty \frac{u-[u]-\frac{1}{2}}{(u+a)^{s+1}} du.$

When $\Re s \ge \frac{1}{2}$, we choose $y = 1$, then

$\zeta(s,a) = a^{-s} + (1+a)^{-s} + \frac{(1+a)^{1-s}}{s-1} - \frac{1}{2}(1+a)^{-s} - s\int_1^\infty \frac{u-[u]-\frac{1}{2}}{(u+a)^{s+1}} du,$

hence

$$\zeta(s,a) \ll a^{-\sigma} + | t |, \quad | t | \ge 1.$$

When $\frac{1}{k} - 1 < \Re s \le \frac{1}{2}$, we choose $y = | t | \ge 1$, then

$\zeta(s,a) \ll a^{-\sigma} + \sum\limits_{1 \le n \le |t|} \frac{1}{(n+a)^\sigma} + | t |^{1-\sigma} + | s\int_{|t|}^\infty \frac{u-[u]-\frac{1}{2}}{(u+a)^{s+1}} du |,$

by partial integrations,

$| s\int_{|t|}^\infty \frac{u-[u]-\frac{1}{2}}{(u+a)^{s+1}} du | \ll | t |^{1-\sigma} + | t |^2 | t |^{-\sigma-1} \ll | t |^{1-\sigma}.$

We have

$\zeta(s,a) \ll a^{-\sigma} + | t | + | t |^{1-\sigma}, \quad \frac{1}{k} - 1 + \tau \le \Re s \le 2, \ | t | \ge 1.$

By (9.6), (9.21) and $1 \le A_1, A_3, A_4, A_5 \ll d_2^k d_3^k$,

$$G(u;s) \ll 1 + | t | + | t |^{1-\sigma}, \qquad \frac{1}{k} - 1 + \tau \le \Re s \le 2, \ | t | \ge 1,$$

the lemma follows.

By Lemma 10.1,

$\sum\limits_{u \le Q} \frac{1}{2\pi i}\int_{2-iT}^{\sigma-iT} u^{-s} G(u;s) \frac{x^{s+1}}{s(s+1)} ds, \ \sum\limits_{u \le Q} \frac{1}{2\pi i}\int_{\sigma+iT}^{2+iT} u^{-s} G(u;s) \frac{x^{s+1}}{s(s+1)} ds \ll x^3 Q^{1-\sigma} T^{-\frac{1}{k}},$

therefore

$$\begin{aligned} A = & \sum_{u \le Q} \frac{1}{2\pi i}\int_{\sigma-iT}^{\sigma+iT} u^{-s} G(u;s) \frac{x^{s+1}}{s(s+1)} ds \\ & + Res_{s=0,1}\Big(\sum_{u \le Q} u^{-s} G(u;s) \frac{x^{s+1}}{s(s+1)}\Big) + O(x^3 Q^{1-\sigma} T^{-\frac{1}{k}}). \end{aligned} \tag{10.2}$$

Write

$B_1 = \sum\limits_{u \le Q} \frac{1}{2\pi i}\int_{\sigma-iT}^{\sigma+iT} u^{-s} G_{(1)}(u,s) \frac{x^{s+1}}{s(s+1)} ds,$

then, by (9.19),

$$B_1 = \sum_{u \le Q} \tfrac{1}{2\pi i} \int_{\sigma - iT}^{\sigma + iT} u^{-s} \sum_{\substack{(d_2 d_3, u)=1 \\ (d_1 d_2 d_3, 2)=1}} h(d_1, d_2, d_3) \tfrac{(u, 2^k d_1^k)^s}{2^{ks} d_1^{ks}} \zeta(s; A_1, d_2^k d_3^k) \tfrac{x^{s+1}}{s(s+1)} ds$$

$$= \sum_{u \le Q} \tfrac{1}{2\pi i} \int_{\sigma - iT}^{\sigma + iT} u^{-s} \sum_{\substack{(d_2 d_3, u)=1 \\ (d_1 d_2 d_3, 2)=1}} h(d_1, d_2, d_3) \tfrac{(u, 2^k d_1^k)^s}{2^{ks} d_1^{ks}} G(s) (d_2 d_3)^{-ks} (e^{\frac{1}{2} i\pi s} \zeta(1 - s; \xi_{d_2^k d_3^k}^{A_1}) -$$

$$e^{\frac{-1}{2} i\pi s} \zeta(1 - s; \xi_{d_2^k d_3^k}^{-A_1})) \tfrac{x^{s+1}}{s(s+1)} ds = B_{11} + B_{12}, \text{ say.}$$

We have

$$\int_{\sigma - iT}^{\sigma + iT} u^{-s} \tfrac{(u, 2^k d_1^k)^s}{2^{ks} d_1^{ks}} G(s) (d_2 d_3)^{-ks} e^{\frac{1}{2} i\pi s} \zeta(1 - s; \xi_{d_2^k d_3^k}^{A_1}) \tfrac{x^{s+1}}{s(s+1)} ds$$

$$= \sum_{m=1}^{\infty} \int_{\sigma - iT}^{\sigma + iT} u^{-s} \tfrac{(u, 2^k d_1^k)^s}{2^{ks} d_1^{ks}} G(s) (d_2 d_3)^{-ks} e^{\frac{1}{2} i\pi s} \xi_{d_2^k d_3^k}^{A_1 m} m^{-(1-s)} \tfrac{x^{s+1}}{s(s+1)} ds$$

$$= e^{\frac{1}{2} i\pi\sigma} x^{\sigma+1} i u^{-\sigma} \tfrac{(u, 2^k d_1^k)^\sigma}{2^{k\sigma} d_1^{k\sigma}} (d_2 d_3)^{-k\sigma} \sum_{m=1}^{\infty} \xi_{d_2^k d_3^k}^{A_1 m} m^{-1+\sigma} \int_{-T}^{T} u^{-it} \tfrac{(u, 2^k d_1^k)^{it}}{2^{kti} d_1^{kti}} G(\sigma + it) (d_2 d_3)^{-kti} e^{-\frac{1}{2}\pi t} m^{it} \times$$

$$\tfrac{x^{it}}{(\sigma + it)(1 + \sigma + it)} dt,$$

and

$$\int_{-T}^{T} u^{-it} \tfrac{(u, 2^k d_1^k)^{it}}{2^{kti} d_1^{kti}} G(\sigma + it) (d_2 d_3)^{-kti} e^{-\frac{1}{2}\pi t} m^{it} \tfrac{x^{it}}{(\sigma + it)(1 + \sigma + it)} dt$$

$$\ll (\int_{1}^{T} + \int_{-T}^{-1}) u^{-it} \tfrac{(u, 2^k d_1^k)^{it}}{2^{kti} d_1^{kti}} G(\sigma + it) (d_2 d_3)^{-kti} e^{-\frac{1}{2}\pi t} m^{it} \tfrac{x^{it}}{(\sigma + it)(1 + \sigma + it)} dt + 1.$$

Let $1 \le T_1 \le T_2 \le 2T_1$. By Lemma 10.1,

$$\int_{T_1}^{T_2} u^{-it} \tfrac{(u, 2^k d_1^k)^{it}}{2^{kti} d_1^{kti}} G(\sigma + it) (d_2 d_3)^{-kti} e^{-\frac{1}{2}\pi t} m^{it} \tfrac{x^{it}}{(\sigma + it)(1 + \sigma + it)} dt$$

$$= -i \int_{T_1}^{T_2} u^{-it} \tfrac{(u, 2^k d_1^k)^{it}}{2^{kti} d_1^{kti}} (2\pi)^{\sigma + it - 1} \sqrt{2\pi} e^{-\frac{\pi}{2} t} t^{1 - \sigma - \frac{1}{2}} e^{-it(\log t - 1)} e^{-\frac{i\pi}{2}(1 - \sigma - \frac{1}{2})} (1 + O(t^{-1})) \times$$

$$(d_2 d_3)^{-kti} e^{-\frac{1}{2}\pi t} m^{it} \tfrac{x^{it}}{(\sigma + it)(1 + \sigma + it)} dt \ll T_1^{-\frac{3}{2} - \sigma},$$

and

$$\int_{T_1}^{T_2} u^{it} \tfrac{(u, 2^k d_1^k)^{-it}}{2^{-kti} d_1^{-kti}} G(\sigma - it) (d_2 d_3)^{kti} e^{\frac{1}{2}\pi t} m^{-it} \tfrac{x^{-it}}{(\sigma - it)(1 + \sigma - it)} dt$$

$$= -i (2\pi)^{\sigma - \frac{1}{2}} e^{\frac{i\pi}{2}(\frac{1}{2} - \sigma)} \int_{T_1}^{T_2} u^{it} \tfrac{(u, 2^k d_1^k)^{-it}}{2^{-kti} d_1^{-kti}} (2\pi)^{-it} t^{1 - \sigma - \frac{1}{2}} e^{it(\log t - 1)} (1 + O(t^{-1})) \times$$

$$(d_2 d_3)^{kti} m^{-it} \tfrac{x^{-it}}{(\sigma - it)(1 + \sigma - it)} dt$$

$$= i (2\pi)^{\sigma - \frac{1}{2}} e^{\frac{i\pi}{2}(\frac{1}{2} - \sigma)} \int_{T_1}^{T_2} u^{it} \tfrac{(u, 2^k d_1^k)^{-it}}{2^{-kti} d_1^{-kti}} (2\pi)^{-it} t^{-\frac{3}{2} - \sigma} e^{it(\log t - 1)} (d_2 d_3)^{kti} m^{-it} x^{-it} dt + O(T_1^{-\frac{3}{2} - \sigma}).$$

Write

$$B = \int_{T_1}^{T_2} u^{it} \tfrac{(u, 2^k d_1^k)^{-it}}{2^{-kti} d_1^{-kti}} (2\pi)^{-it} t^{-\frac{3}{2} - \sigma} e^{it(\log t - 1)} (d_2 d_3)^{kti} m^{-it} x^{-it} dt = \int_{T_1}^{T_2} t^{-\frac{3}{2} - \sigma} e^{iF(t)} dt,$$

where

$$F(t) = t\log u - t\log(u, 2^k d_1^k) + kt\log(2d_1d_2d_3) - t\log(2\pi) + t(\log t - 1) - t\log(mx).$$

$$F''(t) = t^{-1},$$

therefore, by Lemma 4.4 of [7] and by partial integrations,

$\int_{T_1}^{T_2} e^{iF(t)}dt \ll T_1^{\frac{1}{2}}, \quad B \ll T_1^{-1-\sigma}.$

Hence

$$\int_{-T}^{T} u^{-it}\frac{(u, 2^k d_1^k)^{it}}{2^{kti}d_1^{kti}}G(\sigma+it)(d_2d_3)^{-kti}e^{-\frac{1}{2}\pi t}m^{it} \times \frac{x^{it}}{(\sigma+it)(1+\sigma+it)}dt \ll 1,$$

and, by (9.6),

$$B_{11} \ll \sum_{u\leq Q} x^{\sigma+1}u^{-\sigma} \ll x^{\sigma+1}Q^{1-\sigma}.$$

The same proof works for B_{12}, then

$$B_{12} \ll x^{\sigma+1}Q^{1-\sigma},$$

we obtain

$$\sum_{u\leq Q}\frac{1}{2\pi i}\int_{\sigma-iT}^{\sigma+iT} u^{-s}G_{(1)}(u,s)\frac{x^{s+1}}{s(s+1)}ds \ll x^{\sigma+1}Q^{1-\sigma}.$$

Similarly, we deduce

$$\sum_{u\leq Q}\frac{1}{2\pi i}\int_{\sigma-iT}^{\sigma+iT} u^{-s}G_{(j)}(u,s)\frac{x^{s+1}}{s(s+1)}ds \ll x^{\sigma+1}Q^{1-\sigma}, \quad j = 3,4,5.$$

By (9.21),

$$\sum_{u\leq Q}\frac{1}{2\pi i}\int_{\sigma-iT}^{\sigma+iT} u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}ds \ll x^{\sigma+1}Q^{1-\sigma}. \tag{10.3}$$

Let $T \to \infty$, by (10.1), (10.2) and (10.3),

$$S_E(x,Q) = Res_{s=0,1}(\sum_{u\leq Q} u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}) + O(x^{\sigma+1}Q^{1-\sigma}). \tag{10.4}$$

We have

$$Res_{s=1}(\sum_{u\leq Q} u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}) = c_2(Q)x^2, \tag{10.5}$$

where $c_2(Q) = \frac{1}{2}\sum\limits_{u\le Q} u^{-1}Res_{s=1}G(u;s)$, $c_2(Q)$ is independence of x.

We need the value $\zeta(0;a,k) = \frac{1}{2} - \frac{a}{k},\ 1 \le a \le k$.

$$Res_{s=0}(\sum_{u\le Q} u^{-s}G(u;s)\frac{x^{s+1}}{s(s+1)}) = x\sum_{u\le Q} G(u;0). \tag{10.6}$$

LEMMA 10.2 We have

$\sum\limits_{u\le Q} G_{(1)}(u,0) = c_{31}Q + O(Q^{\frac{1}{k}+\varepsilon}),\quad \sum\limits_{u\le Q} G_{(3)}(u,0) = c_{33}Q + O(Q^{\frac{1}{k}+\varepsilon}),$
$\sum\limits_{u\le Q} G_{(4)}(u,0) = c_{34}Q + O(Q^{\frac{1}{k}+\varepsilon}),\quad \sum\limits_{u\le Q} G_{(5)}(u,0) = c_{35}Q + O(Q^{\frac{1}{k}+\varepsilon}),$
where $c_{31}, c_{33}, c_{34}, c_{35}$ are independence of Q.

Proof. We only consider the case $(d_i, d_l) = 1, i \ne l$.

$$\begin{aligned}\sum_{u\le Q} G_{(1)}(u,0) &= \sum_{u\le Q}\ \sum_{\substack{(d_2d_3,u)=1\\(d_1d_2d_3,2)=1}} h(d_1,d_2,d_3)\zeta(0;A_1,d_2^kd_3^k)\\ &= \sum_{u\le Q}\ \sum_{\substack{(d_2d_3,u)=1\\(d_1d_2d_3,2)=1}} h(d_1,d_2,d_3)(\tfrac{1}{2} - A_1d_2^{-k}d_3^{-k}).\end{aligned} \tag{10.7}$$

We have

$\sum\limits_{\substack{u\le Q\\(m,u)=1}} 1 = Q\sum\limits_{t|m}\mu(t)t^{-1} + O(\tau(m)),$
then

$\sum\limits_{u\le Q}\ \sum\limits_{\substack{(d_2d_3,u)=1\\(d_1d_2d_3,2)=1}} h(d_1,d_2,d_3)\frac{1}{2} = \frac{1}{2}\sum\limits_{(d_1d_2d_3,2)=1} h(d_1,d_2,d_3)\sum\limits_{\substack{u\le Q\\(d_2d_3,u)=1}} 1$
$= \frac{1}{2}\sum\limits_{(d_1d_2d_3,2)=1} h(d_1,d_2,d_3)(Q\sum\limits_{t|d_2d_3}\mu(t)t^{-1} + O(\tau(d_2d_3))),$
therefore, by (9.6),

$$\sum_{u\le Q}\ \sum_{\substack{(d_2d_3,u)=1\\(d_1d_2d_3,2)=1}} h(d_1,d_2,d_3)\frac{1}{2} = \frac{1}{2}Q\sum_{(d_1d_2d_3,2)=1} h(d_1,d_2,d_3)\sum_{t|d_2d_3}\mu(t)t^{-1} + O(1). \tag{10.8}$$

Write

$C = \sum\limits_{\substack{u\le Q\\(d_2d_3,u)=1}} A_1 = \sum\limits_{\substack{u\le Q\\(d_2d_3,u)=1}} \{-b_1d_3^k\overline{d_3^k} + b_2d_2^k\overline{d_2^k}\}',$
where $A_1 = \{-b_1d_3^k\overline{d_3^k} + b_2d_2^k\overline{d_2^k}\}'$ means $1 \le A_1 \le d_2^kd_3^k, (A_1, d_2d_3) = 1$.

Let $(u, 2^kd_1^k) = j,\ u = ju_1$, then, by (9.8),(9.10) and (9.10),

$$2^kd_1^ku_1b_1 \equiv 1(mod\ d_2^k),\ 2^kd_1^ku_1b_2 \equiv 1(mod\ d_3^k),$$

and

$$C = \sum_{j_1=1}^{d_2^k}{}' \sum_{j_2=1}^{d_3^k}{}' \{-j_1 d_3^k \overline{d_3^k} + j_2 d_2^k \overline{d_2^k}\}' \sum_{j \mid 2^k d_1^k} D,$$

where

$$D = \#\{u_1 : u_1 \le Qj^{-1}, (u_1, 2^k d_1^k j^{-1}) = 1, 2^k d_1^k u_1 j_1 \equiv 1 (mod\ d_2^k), \\ 2^k d_1^k u_1 j_2 \equiv 1 (mod\ d_3^k)\}.$$

$$D = \sum_{t \mid 2^k d_1^k j^{-1}} \mu(t) \sum_{\substack{u \le Qj^{-1}t^{-1} \\ u 2^k d_1^k t j_1 \equiv 1 (mod\ d_2^k), u 2^k d_1^k t j_2 \equiv 1 (mod\ d_3^k)}} 1 = \sum_{t \mid 2^k d_1^k j^{-1}} \mu(t) \sum_{\substack{u \le Qj^{-1}t^{-1} \\ u \equiv J (mod\ d_2^k d_3^k)}} 1,$$

where

$$J \equiv \overline{2^k d_1^k t}\, \overline{j_1} d_3^k \overline{d_3^k} + \overline{2^k d_1^k t}\, \overline{j_2} d_2^k \overline{d_2^k}\ (mod\ d_2^k d_3^k), \text{and } 1 \le J \le d_2^k d_3^k. \tag{10.9}$$

We have

$$\sum_{\substack{u \le Qj^{-1}t^{-1} \\ u \equiv J (mod\ d_2^k d_3^k)}} 1 = \begin{cases} Qj^{-1}t^{-1}d_2^{-k}d_3^{-k} + O((Qj^{-1}t^{-1}d_2^{-k}d_3^{-k})^{\frac{1}{k}+\varepsilon}), & \text{if } d_2^k d_3^k \le Qj^{-1}t^{-1}, \\ 1, & \text{if } d_2^k d_3^k > Qj^{-1}t^{-1} \ge J, \\ 0, & \text{if } d_2^k d_3^k \ge J > Qj^{-1}t^{-1}. \end{cases}$$

Therefore

$$\sum_{u \le Q} \sum_{\substack{(d_2 d_3, u)=1 \\ (d_1 d_2 d_3, 2)=1}} h(d_1, d_2, d_3) A_1 d_2^{-k} d_3^{-k}$$

$$= \sum_{(d_1 d_2 d_3, 2)=1} h(d_1, d_2, d_3) d_2^{-k} d_3^{-k} \sum_{j_1=1}^{d_2^k}{}' \sum_{j_2=1}^{d_3^k}{}' \{-j_1 d_3^k \overline{d_3^k} + j_2 d_2^k \overline{d_2^k}\}' \sum_{j \mid 2^k d_1^k} \sum_{t \mid 2^k d_1^k j^{-1}} \mu(t) \sum_{\substack{u \le Qj^{-1}t^{-1} \\ u \equiv J (mod\ d_2^k d_3^k)}} 1$$

$$= \sum_{(d_1,2)=1} \sum_{j \mid 2^k d_1^k} \sum_{t \mid 2^k d_1^k j^{-1}} \mu(t) \sum_{d_2^k d_3^k \le Qj^{-1}t^{-1}} h(d_1, d_2, d_3) d_2^{-k} d_3^{-k} \times$$

$$\sum_{j_1=1}^{d_2^k}{}' \sum_{j_2=1}^{d_3^k}{}' \{-j_1 d_3^k \overline{d_3^k} + j_2 d_2^k \overline{d_2^k}\}' (Qj^{-1}t^{-1}d_2^{-k}d_3^{-k} + O((Qj^{-1}t^{-1}d_2^{-k}d_3^{-k})^{\frac{1}{k}+\varepsilon}))$$

$$+ \sum_{(d_1,2)=1} \sum_{j \mid 2^k d_1^k} \sum_{t \mid 2^k d_1^k j^{-1}} \mu(t) \sum_{d_2^k d_3^k > Qj^{-1}t^{-1}} h(d_1, d_2, d_3) d_2^{-k} d_3^{-k} \sum_{j_1=1}^{d_2^k}{}' \sum_{\substack{j_2=1 \\ 1 \le J \le Qj^{-1}t^{-1}}}^{d_3^k}{}' \{-j_1 d_3^k \overline{d_3^k} +$$

$$j_2 d_2^k \overline{d_2^k}\}'.$$

By (9.6),

$$\sum_{(d_1,2)=1} \sum_{j \mid 2^k d_1^k} \sum_{t \mid 2^k d_1^k j^{-1}} \mu(t) \sum_{\substack{(d_2 d_3, 2)=1 \\ d_2^k d_3^k \le Qj^{-1}t^{-1}}} h(d_1, d_2, d_3) d_2^{-k} d_3^{-k} \times$$

$$\sum_{j_1=1}^{d_2^k}{}' \sum_{j_2=1}^{d_3^k}{}' \{-j_1 d_3^k \overline{d_3^k} + j_2 d_2^k \overline{d_2^k}\}' (Qj^{-1}t^{-1}d_2^{-k}d_3^{-k} + O((Qj^{-1}t^{-1}d_2^{-k}d_3^{-k})^{\frac{1}{k}+\varepsilon}))$$

$$= Q \sum_{(d_1 d_2 d_3, 2)=1} \sum_{j | 2^k d_1^k} \sum_{t | 2^k d_1^k j^{-1}} \mu(t) j^{-1} t^{-1} h(d_1, d_2, d_3) d_2^{-2k} d_3^{-2k} \sum_{a=1}^{d_2^k d_3^k}{}' a + O(Q^{\frac{1}{k}+\varepsilon}),$$

we note that : if $tj > Q$, this will produce the error term $O(1)$, therefore the above formula is valid.

By (9.6) and (10.9),

$$\sum_{(d_1,2)=1} \sum_{j | 2^k d_1^k} \sum_{t | 2^k d_1^k j^{-1}} \mu(t) \sum_{d_2^k d_3^k > Q j^{-1} t^{-1}} h(d_1, d_2, d_3) d_2^{-k} d_3^{-k} \times$$
$$\sum_{j_1=1}^{d_2^k}{}' \sum_{\substack{j_2=1 \\ 1 \le J \le Q j^{-1} t^{-1}}}^{d_3^k} {}' \{ -j_1 d_3^k \overline{d_3^k} + j_2 d_2^k \overline{d_2^k} \}'$$
$$\ll \sum_{(d_1,2)=1} \sum_{j | 2^k d_1^k} \sum_{t | 2^k d_1^k j^{-1}} \mu(t)^2 \sum_{d_2^k d_3^k > Q j^{-1} t^{-1}} h(d_1, d_2, d_3) d_2^{-k} d_3^{-k} d_2^k d_3^k Q j^{-1} t^{-1}$$
$$\ll Q \sum_{(d_1,2)=1} \sum_{j | 2^k d_1^k} \sum_{t | 2^k d_1^k j^{-1}} \mu(t)^2 j^{-1} t^{-1} \sum_{d_2^k d_3^k > Q j^{-1} t^{-1}} h(d_1, d_2, d_3) \ll Q^{\frac{1}{k}+\varepsilon}.$$

Hence, by (10.7) and (10.8),

$$\sum_{u \le Q} G_{(1)}(u, 0) = c_{31} Q + O(Q^{\frac{1}{k}+\varepsilon}),$$

where c_{31} is independence of Q.

Similarly, by the same method, we can obtain

$$\sum_{u \le Q} G_{(3)}(u, 0) = c_{33} Q + O(Q^{\frac{1}{k}+\varepsilon}), \sum_{u \le Q} G_{(4)}(u, 0) = c_{34} Q + O(Q^{\frac{1}{k}+\varepsilon}),$$
$$\sum_{u \le Q} G_{(5)}(u, 0) = c_{35} Q + O(Q^{\frac{1}{k}+\varepsilon}),$$

where c_{33}, c_{34}, c_{35} are independence of Q, the lemma follows.

Consequently, for $\frac{1}{k} - 1 + \tau \le \sigma < 0$, by Lemma 10.2, (9.21), (10.4), (10.5) and (10.6),

$$S_E(x, Q) = c_2(Q) x^2 + c_3 x Q + O(x^{\sigma+1} Q^{1-\sigma} + x Q^{\frac{1}{k}+\varepsilon}),$$

by choosing $\sigma = \frac{1}{k} - 1 + \tau < 0$, we deduce

$$S_E(x, Q) = c_2(Q) x^2 + c_3 x Q + O(x^{\frac{1}{k}+\tau} Q^{2-\frac{1}{k}-\tau} + x Q^{\frac{1}{k}+\varepsilon}), \tag{10.10}$$

where $c_2(Q)$ is independence of x , and c_3 is independence of x, Q.

11 Proof of the Theorem

When $k = 2$, the Theorem follows from Lemma 5.2. We suppose that $k > 2$.

We also note that $R = x^{\frac{1}{2}+\tau}$. By (4.33), (5.6), (6.4), (7.2) and (10.9),

$$S_1(x,Q) = 2c_2(Q)x^2 + 2c_3xQ + [Q]\sum_{n\le x}\mu_k(n)\mu_k(n+1) + O(x^{\frac{1}{k}+\tau}Q^{2-\frac{1}{k}-\tau} + x^{\frac{3}{2}+\frac{1}{2k}+2\tau}) \tag{11.1}$$

By Theorem 1 of [4],

$$\sum_{n\le x}\mu_k(n)\mu_k(n+1) = \varrho x + O(x^{\frac{2}{k+1}}),$$

where ϱ is defined by (4.17).

Hence, by (1.5), (1.6), (2.32), (3.4) and (11.1),

$$Y_k(x,Q) = c_4(Q)x^2 + c_5xQ + O(x^{\frac{1}{k}+\tau}Q^{2-\frac{1}{k}-\tau} + x^{\frac{3}{2}+\frac{1}{2k}+2\tau}),$$

and

$$c_4(Q)x^2 + c_5xQ + O(x^{\frac{1}{k}+\tau}Q^{2-\frac{1}{k}-\tau} + x^{\frac{3}{2}+\frac{1}{2k}+2\tau}) \ll x^{\frac{2}{k}+\varepsilon}Q^{2-\frac{2}{k}} + x^{\frac{4}{k+1}+\varepsilon}, \tag{11.2}$$

where $c_4(Q)$ is independence of x , and c_5 is independence of x, Q. (11.2) is valid for all $1 < Q \le x$.

For fixed Q, we divide both sides of (11.2) by x^2 and let $x \to \infty$, we deduce $c_4(Q) = 0$ for all $Q > 1$.

hence

$$Y_k(x,Q) = c_5xQ + O(x^{\frac{1}{k}+\tau}Q^{2-\frac{1}{k}-\tau} + x^{\frac{3}{2}+\frac{1}{2k}+2\tau}),$$

and

$$c_5xQ + O(x^{\frac{1}{k}+\tau}Q^{2-\frac{1}{k}-\tau} + x^{\frac{3}{2}+\frac{1}{2k}+2\tau}) \ll x^{\frac{2}{k}+\varepsilon}Q^{2-\frac{2}{k}} + x^{\frac{4}{k+1}+\varepsilon}, \tag{11.3}$$

We divide both sides of (11.3) by xQ , by choosing $Q = x^{\frac{5}{6}}$ and let $x \to \infty$, we obtain $c_5 = 0$.

Consequently,

$$Y_k(x,Q) = O(x^{\frac{1}{k}+\tau}Q^{2-\frac{1}{k}-\tau} + x^{\frac{3}{2}+\frac{1}{2k}+2\tau}),$$

since τ is a sufficiently small positive number, the Theorem follows at once.

References

[1] J. Brüdern, A. Granville, A. Perelli, R. C. Vaughan & T. D. Wooley, On the exponential sum over $k-$free numbers, Phil. Trans. R. Soc. London A 356 (1998)739-761.

[2] J. Brüdern, A. Perelli & T. D. Wooley, Twins of $k-$free numbers and their exponential sum, Michigan Math.J. 47 (2000) 173-190.

[3] T.Estermann, On the representations of a number as the sum of two products, Proc. London Math. Soc. (2) 31(1930) 123-133.

[4] Z.Z.Meng, Some new results on $k-$free numbers, J.Number Theory, 121(2006) 45-66.

[5] C.D.Pan.& C.B.Pan, The foundation of analytic number theory(in Chinese),Beijing, Science Press, 1997.

[6] E.C.Titchmarsh, The theory of functions, Oxford University Press,1952.

[7] E.C.Titchmarsh, The theory of the Riemann zeta-function, 2nd edn, revised by D.R.Heath-Brown(Oxford University Press,1986).

[8] R. C. Vaughan, The Hardy-Littlewood method,2nd edn,Cambridge Tracts in Mathematics 125(Cambridge University Press,1997).

[9] R. C. Vaughan, A variance for $k-$free numbers in arithmetic progressions, Proc. London Math. Soc. (3) 91(2005) 573-597.

School of Mathematical Sciences

Peking University

Beijing, 100871, P.R.China

E-mail: mzzh@math.pku.edu.cn